

# Exhibit 295



# New insights on occupational exposure and bladder cancer risk: a pooled analysis of two Italian case–control studies

Veronica Sciannameo<sup>1,2</sup> · Angela Carta<sup>3,4</sup> · Angelo d’Errico<sup>1</sup> · Maria Teresa Giraudo<sup>5</sup> · Francesca Fasanelli<sup>6</sup> · Cecilia Arici<sup>4,10</sup> · Milena Maule<sup>6</sup> · Paolo Carnà<sup>1</sup> · Paolo Destefanis<sup>7</sup> · Luigi Rolle<sup>7</sup> · Paolo Gontero<sup>7</sup> · Giovanni Casetta<sup>7</sup> · Andrea Zitella<sup>7</sup> · Giuseppina Cucchiarale<sup>8</sup> · Paolo Vineis<sup>9</sup> · Stefano Porru<sup>4,10</sup> · Carlotta Sacerdote<sup>6</sup> · Fulvio Ricceri<sup>1,2</sup>

Received: 10 January 2018 / Accepted: 24 November 2018 / Published online: 30 November 2018  
© Springer-Verlag GmbH Germany, part of Springer Nature 2018

## Abstract

**Purpose** The main risk factor for bladder cancer (BC) is cigarette smoking, but also occupational exposure to carcinogens is relevant, causing about 4–10% of BC. We aimed at investigating the association between BC risk, occupations held in the past and exposure to occupational carcinogens, also assessing whether these associations were influenced by tumour grade.

**Methods** We pooled data from two Italian case–control studies on male BC, analyzing 893 cases and 978 controls. Occupations were classified using the International Standard Classification of Occupations and exposure to carcinogens was assigned using a validated Job Exposure Matrix. Logistic regression approach was used as well as a semi-Bayesian model, based on a priori information on exposure.

**Results** A significantly increased BC risk was found for chemical engineering technicians, postmen, and lathe operators, but only, for the latter, the association remained significant after Bayesian control for type I error. Among carcinogens, cadmium and trichloroethylene were associated with BC. When analyzing data by grade, exposure to these carcinogens was associated with low-grade BC only.

**Conclusions** Our results suggest that monitoring workplaces to prevent exposure to carcinogenic agents is still an important task, which should be still given adequate importance in public health.

**Keywords** Bladder cancer · Occupational exposure · Carcinogens · Case–control study · Bayesian methods

---

Carlotta Sacerdote and Fulvio Ricceri are joint last authors.

**Electronic supplementary material** The online version of this article (<https://doi.org/10.1007/s00420-018-1388-2>) contains supplementary material, which is available to authorized users.

✉ Fulvio Ricceri  
[fulvio.ricceri@unito.it](mailto:fulvio.ricceri@unito.it)

<sup>1</sup> Unit of Epidemiology, Regional Health Service, ASLTO3, Via Sabaudia 164, 10095 Grugliasco, TO, Italy

<sup>2</sup> Department of Biological and Clinical Sciences, University of Turin, 10043 Orbassano, TO, Italy

<sup>3</sup> Department of Medical-Surgical Specialties, Radiological Sciences and Public Health, University of Brescia, 25123 Brescia, Italy

<sup>4</sup> University Research Center Integrated Models for Prevention and Protection in Environmental and Occupational Health, University of Brescia, 25123 Brescia, Italy

## Introduction

Bladder cancer (BC) is the most common neoplasm of the genitourinary tract and it is the ninth most diffused cancer in the world, with 430,000 new cases every year (Stewart and Wild 2014). Among people of Caucasian origin, about

<sup>5</sup> Department of Mathematics “G. Peano”, University of Turin, 10123 Turin, Italy

<sup>6</sup> Unit of Cancer Epidemiology, CPO Piedmont, Department of Medical Sciences, Città della Salute e della Scienza Hospital, University of Turin, 10126 Turin, Italy

<sup>7</sup> Unit of Urology U, Città della Salute e della Scienza Hospital, University of Turin, 10126 Turin, Italy

<sup>8</sup> Unit of Urology I, Cellini Clinic, 10126 Turin, Italy

<sup>9</sup> Imperial College London, London W2 1PG, UK

<sup>10</sup> Department of Diagnostics and Public Health, University of Verona, 37134 Verona, Italy

92–99% of BCs are urothelial and most of them develop in the bladder's wall, less frequently in the trigone, in the urothelial orifices, in the dome, and in the bladder neck (Schottenfeld and Fraumeni 2006).

In Italy, BC represents about 11% of the total cancer diagnoses among men and 3% among women (AIOM AIR-TUM 2016), and data from cancer registries (2005–2009) reveal an age-standardized rate equal to 45.4 cases per 100,000 in men and 8.2 cases per 100,000 in women (ITACAN v2.0 2018).

Risk factors for bladder cancer are mostly known; among them, cigarette smoking is the leading causal factor, causing more than 40% men's BC (Van Osch et al. 2016). Other risk factors are: urinary system chronic infections (Negri and La Vecchia 2001), ionizing radiations treatments (Weiss et al. 1994), disinfection byproducts in water (Hrudey et al. 2015), diet poor in fruit and vegetables (Sacerdote et al. 2007), and the presence of arsenic in drinking water (Saint-Jacques et al. 2014). An increased risk of bladder cancer is also observed among atomic bomb survivors and workers of nuclear installations (McGeoghegan and Binks 2000).

Furthermore, in industrialized countries, the second most important cause of BC is the exposure to occupational carcinogens, causing about 4–10% bladder tumour cases (Kogevinas et al. 1998, 2003). From the '50s on, a series of studies have detected about 40 occupations associated with an increased risk of BC (Kogevinas et al. 1998, 2003; Case and Hosker 1954; Case et al. 1954; Samanic et al. 2008; Purdue et al. 2015). Among those, the International Agency for Research on Cancer (IARC) has classified several occupations as carcinogenic to humans—group 1 (painters, workers in the rubber industry, in aluminum production, and in the production of the dyes auramine and magenta), or as probably carcinogenic to humans—group 2A (hairdressers or barbers), or as possibly carcinogenic to humans—group 2B (workers employed in dry cleaning, in printing processes, and in textile manufacturing). Moreover, an increased risk is present in workers exposed to aromatic amines (Vineis and Pirastu 1997) and polycyclic aromatic hydrocarbons (Boffetta and Silverman 2001; Rota et al. 2014).

To our knowledge, all these studies investigated solely either occupations or exposure to carcinogens; moreover, the investigation of several determinants together could bring to spurious through statistically significant results, due to the risk of type I error.

For this reason, the objectives of our study are:

1. to analyze the association between past employment and BC risk, using a Job Exposure Matrix (JEM), to use a priori information about the supposed presence of carcinogens in the working environment;
2. to analyze the relation between exposure to selected occupational carcinogens and BC risk;

3. to assess whether the association with occupational exposure to carcinogens differs by BC tumour grade.

## Materials and methods

### Subjects

For this study, we pooled data from two existing case–control studies on male BC conducted in the metropolitan areas of Turin and Brescia, two highly industrialized cities located in Northern Italy.

Details of the two studies can be found elsewhere (Ricceri et al. 2010; Sacerdote et al. 2013; Porru et al. 1996). Within the Brescia study, Porru et al. (2014) have already examined the interaction of previous exposure to PAHs and aromatic amines with the presence of genetic polymorphisms of metabolic and DNA repair enzymes on BC risk. Briefly, in Turin, 603 cases and 681 hospital controls were recruited between 1993 and 2012. Cases were patients aged between 40 and 74 years, with a new diagnosis of BC (ICD-IX 188.XX), histologically confirmed and diagnosed at S. Giovanni Battista hospital and Cellini clinic in Turin. Controls were patients aged between 40 and 74 years, hospitalized in S. Giovanni Battista in Turin, not affected by neoplastic, metabolic, urologic, or smoking-related diseases. They were recruited in general medicine, otolaryngology, orthopedics, and cardiology departments, with each disease present in no more than 10% of controls. Response rate was higher than 97% both in cases and controls. In Brescia 295 cases and 302 controls were recruited in 1992 and 1993 and from 1997 to 2000. Cases were patients aged between 20 and 80 years with a new diagnosis of BC, histologically confirmed and diagnosed in the urology departments of the two main hospitals of the city. Controls were selected in the same hospitals among patients hospitalized for urological non-neoplastic diseases and they were frequency-matched to cases by age ( $\pm 5$  years), period (1992–1993 and 1997–2000), and hospital of admission. Response rate was 93% in cases and 97% in controls.

Tumour histology, site, TNM classification, and grading of the primary tumour were assessed through inspection of the patients' pathology and clinical records, in collaboration with the urologists. Tumour stage and grade were determined according to the European Association of Urology Guidelines on Non-Muscle-invasive Urothelial Carcinoma of the Bladder (Babjuk et al. 2013).

### Occupational exposures

In both studies, all patients answered a detailed questionnaire about their working history and lifestyle, including smoking habit. We classified occupations based on the

International Standard Classification of Occupations (ISCO-68) (International Labour Office 1968), using the first four digits, as usually done in similar contexts.

In agreement with the literature, we considered a latency period from exposure to BC of 10 years (Rushton et al. 2008). It means that we assumed that a carcinogen exposure should have occurred at least 10 years before the onset of BC to be involved in the pathogenetic mechanism of that BC. Therefore, only jobs and occupational exposures occurring more than 10 years before BC for cases and before recruitment for controls have been taken into account. For this reason, young subjects that had been working for less than 10 years were excluded from the analyses ( $n = 10$ , 0.53% of the overall recruited subjects), leaving a final sample composed of 602 cases and 680 controls in the Turin study and 291 cases and 298 controls in the Brescia study.

### Carcinogenic agents

The IARC has identified five agents that are possibly involved in the carcinogenic process of BC or tumours of the urinary tract and that are classified as certainly carcinogenic for humans (IARC—group 1) (IARC 2012): arsenic (AS), benzo[a]pyrene (BAP), welding fumes (WELD), aromatic amines (AA: 4-aminobiphenyl, benzidine, 2-naphthylamine, Ortho-Toluidine), and chlorinated hydrocarbon solvents (CHC: trichloroethylene, 1,2-dichloropropane). Moreover, other 24 probable (IARC—group 2A) or possible (IARC—group 2B) carcinogens are suspected risk factors for BC incidence and/or for other cancers of the genitourinary system: aliphatic and alicyclic hydrocarbon solvents (ALHC), diesel engine exhausts (DEEX), aromatic hydrocarbon solvents (ARHC), benzene (BENZ), cadmium (CD), chromium (CR), detergents (DET), fungicides (FUNG), gasoline (GASO), gasoline engine exhaust (GEEX), herbicides (HERB), insecticides (INSC), ionizing radiation (IRAD), iron (FE), lead (PB), low-frequency magnetic fields (MF), methylene chloride (MCH), nickel (NI), oil mist (OIL), other organic solvents (OSOL), perchloroethylene (PER), toluene (TOLU), trichloroethane (TCE), and trichloroethylene (TRI) (IARC 2012; Supplementary Table 1).

For all agents except AA, the exposure was assigned using a Finnish Job Exposure Matrix (FINJEM) (Kauppinen et al. 2009). Exposure to AA, that was absent in the above-cited JEM, was assigned by one of the authors, expert in industrial hygiene (A. d'E.), based on information extracted from different occupational exposure databases (Boiano and Hull 2001; Falcone et al. 2013) and from the last IARC monograph updating exposure to aromatic amines (IARC 2012), with consensus obtained from all the research group. The JEM intensity and probability of exposure measured during years 1960–1984 were used, since it was assumed that, in the study population, most of the relevant exposures

occurred before mid-80's, given the collection period of the subjects and the long latency of BC.

### Descriptive analyses

Data were described using means and standard deviations or frequencies and percentages, for continuous or categorical variables, respectively. Age differences between cases and controls were tested using the Wilcoxon rank-sum test, after ascertainment of non-normality of the distribution through Kolmogorov–Smirnov test, while differences between categorical variables were tested using the Chi-square test.

All statistical analyses were performed using SAS V 9.3 and R software V 3.3.1 and a two-tailed  $p$  value less than 0.05 was considered statistically significant.

### Analyses on occupations

Occupational exposure was analyzed using two different approaches. First, we used a dichotomous variable for each profession, indicating if a subject had ever been employed in that occupation for at least 2 years in his life (even if not continuously); second, we considered another dichotomous variable for each profession, examining the longest occupation held during subject's working life. Only occupations with at least five cases were considered for the analysis.

In each of the two approaches, odds ratios (ORs) and 95% confidence intervals (95% CI) were obtained from multivariable logistic regression models, adjusted for age, smoking status, intensity of smoking, and study (Turin or Brescia). A semi-Bayesian hierarchical model was employed to control for multiple comparisons and to improve the plausibility and stability of estimates [theoretical details of this model can be found in Witte et al. (1994), Greenland (1994), and Corbin et al. (2012)].

### Analyses on occupational exposure to carcinogens

To estimate the risk associated with exposure to carcinogenic agents, ORs and 95% CI were obtained using multivariable logistic regression models, adjusted for age, smoking status, intensity of smoking, and study (Turin or Brescia), considering two types of exposure. First, a never/ever (for at least 2 years) exposure was considered; second, a cumulative exposure was considered, as done in d'Errico et al. (2009), categorizing the probability of exposure to the agent in each profession (from JEM data) into “low” (less than 10% of probability to be exposed), “medium” (between 10 and 50%), and “high” (more than 50%), and the level of exposure, derived from the JEM, into “low”, “medium”, and “high”, using tertiles of each agent distribution as cut-offs. A weight, taking into account intensity and probability of exposure, was

assigned to the different combinations of probability and exposure levels, as shown in Supplementary Table 2. The cumulative exposure of each subject was then obtained multiplying the weight for the duration in years of each job, and summing up this product across the whole subject’s working life. Finally, the cumulative exposure was categorized into three levels: “no exposure”, if the values of exposure were zero, “low exposure”, if the values were less than the median exposure in controls; “high exposure” otherwise.

Since, from exploratory analyses, heterogeneity of the associations by tumour grade was observed for some carcinogens, analyses by grading were also conducted.

To control for the risk of first type error due to multiple comparisons, the False Discovery Rates (FDR) technique was applied (Benjamini and Hochberg 1995).

## Results

In Table 1, results from descriptive analyses are presented separately for each study. The study population was composed of 589 subjects (31.5% of the total sample; 291 cases and 298 controls) from Brescia and 1282 subjects (68.5% of the total sample; 602 cases and 680 controls) from Turin. In the Turin study population, a significant difference in age

**Table 1** Descriptive analyses results

|  | Turin                      |                            | Brescia                    |                            |
|--|----------------------------|----------------------------|----------------------------|----------------------------|
|  | Cases (n=602)              | Controls (n=680)           | Cases (n=291)              | Controls (n=298)           |
|  | Mean (SD) or frequency (%) | Mean (SD) or frequency (%) | Mean (SD) or frequency (%) | Mean (SD) or frequency (%) |
| Age, years                                       | 63.6 (7.8)                 | 59.9 (9.7)                 | 63.7 (10.2)                | 63.7 (10.3)                |
| p value  | <0.0001                    |                            | 0.67                       |                            |
| Smoking status                                   |                            |                            |                            |                            |
| Never smoker                                     | 49 (8.1%)                  | 175 (26.6%)                | 23 (7.9%)                  | 79 (26.5%)                 |
| Former smoker                                    | 289 (48.0%)                | 316 (48.1%)                | 136 (46.7%)                | 139 (46.6%)                |
| Current smoker                                   | 264 (43.8%)                | 166 (25.3%)                | 132 (45.4%)                | 80 (26.8%)                 |
| p value  | <0.0001                    |                            | <0.0001                    |                            |
| Occupation                                       |                            |                            |                            |                            |
| Professional, technical and related workers      | 80 (13.3%)                 | 99 (14.6%)                 | 17 (5.8%)                  | 24 (8.1%)                  |
| Administrative and managerial workers            | 29 (4.8%)                  | 30 (4.4%)                  | 7 (2.4%)                   | 5 (1.7%)                   |
| Clerical and related workers                     | 53 (8.8%)                  | 70 (10.3%)                 | 23 (7.9%)                  | 24 (8.0%)                  |
| Sales workers                                    | 44 (7.3%)                  | 36 (5.3%)                  | 25 (8.6%)                  | 34 (11.4%)                 |
| Service workers                                  | 39 (6.5%)                  | 42 (6.2%)                  | 13 (4.5%)                  | 14 (4.7%)                  |
| Agricultural, animal husbandry, forestry workers | 9 (1.5%)                   | 9 (1.3%)                   | 34 (11.7%)                 | 31 (10.4%)                 |
| Production and related workers, transport        | 348 (57.8%)                | 394 (57.9%)                | 172 (59.1%)                | 166 (55.7%)                |
| p value  | 0.58                       |                            | 0.92                       |                            |
| Tumour type                                      |                            |                            |                            |                            |
| Not invasive                                     | 520 (86.4%)                |                            | 243 (83.5%)                |                            |
| Invasive   | 70 (11.6%)                 |                            | 48 (16.5%)                 |                            |
| Missing  | 12 (2.0%)                  |                            | 0 (0.0%)                   |                            |
| Tumour grade                                     |                            |                            |                            |                            |
| Low  | 364 (60.5%)                |                            | 185 (63.6%)                |                            |
| High   | 238 (39.5%)                |                            | 100 (34.4%)                |                            |
| Missing  | 0 (0.0%)                   |                            | 6 (2.0%)                   |                            |
| Tumour T   |                            |                            |                            |                            |
| In situ  | 9 (1.5%)                   |                            | 14 (4.8%)                  |                            |
| Ta   | 342 (56.7%)                |                            | 70 (24.1%)                 |                            |
| T1   | 170 (28.2%)                |                            | 159 (54.6%)                |                            |
| T2 or more                                       | 70 (11.6%)                 |                            | 48 (16.5%)                 |                            |
| Missing  | 11 (2.0%)                  |                            | 0 (0.00%)                  |                            |

p values were computed performing Wilcoxon or Chi-squared tests

by case status was found (cases: mean 63.6 years, standard deviation SD 7.8; controls: mean 59.9 years, SD 9.7;  $p < 0.0001$ ), while no difference was found in the Brescia study, due to the matching (cases: mean 63.7 years, SD 10.2; controls: mean 63.7 years, SD 10.3,  $p = 0.67$ ). Both studies revealed a significant difference in smoking status ( $p < 0.0001$ ), with higher percentages of smokers among cases (43.8% in Turin; 45.4% in Brescia) than among controls (25.3% in Turin; 26.8% in Brescia). Interestingly, only a small percentage of never smokers were found among cases both in Turin (8.1%) and in Brescia (7.9%).

Results from Chi-square tests showed that cases' and controls' distributions of the occupations longest held in life, categorized in major groups through the first digit of ISCO, were not different ( $p$  value = 0.58, in Turin, and 0.92, in Brescia).

In Turin, 86.4% ( $n = 520$ ) BCs were not invasive, 60.5% ( $n = 364$ ) were low-grade, and 56.7% ( $n = 342$ ) were Ta tumours. In Brescia, 83.5% ( $n = 243$ ) BCs were not invasive, 63.6% ( $n = 185$ ) were low grade, and 24.1% ( $n = 70$ ) were Ta tumours.

### Analyses on occupations

Using the first approach (occupations held for at least 2 years), a list of 128 occupations (described by the first four ISCO-68 digits) were analyzed. All results from the logistic models are presented in Supplementary Table 3 (fourth column). Significantly increased risks were observed for chemical engineering technicians (ISCO-68 0361), OR 12.65 (95% CI 1.30–107.12), for postmen (ISCO-68 3703), OR 5.48 (95% CI 1.12–26.83), for tool and die makers (ISCO-68 8322), OR 6.03 (95% CI 1.19–30.63), and for lathe operators (ISCO-68 8342), OR 1.77 (95% CI 1.09–2.89) (Table 2). Interestingly, there were some occupations that did not have controls, but pertained to an elevated number of cases: rubber and plastic workers (ISCO-68 9019) and spray painters (ISCO-68 9393), with, respectively, 7 and 9 cases and 0 controls for both. In this case, it was not possible to estimate an OR through a regression model, but there was a suggestion of a possible association with BC ( $p$  value for Fisher's exact test: 0.04 and 0.002 respectively). Moreover, when considering all occupations involved in rubber production

as a whole (ISCO-68: 9012, 9013, 9014 and 9019), we observed an increased risk of developing BC (OR 3.25, 95% CI 1.07–9.83).

Performing the Semi-Bayesian second stage of analysis, to control for false-positive results, we found that only the excess risk for lathe operators (ISCO-68 8342) maintained its statistical significance (Supplementary Table 3).

Using the second approach for the occupational exposure analysis (longest occupations performed in life), a list of 50 occupations (described by the first four ISCO-68 digits) were analyzed. Results are reported in Supplementary Table 4, showing again an increased risk of BC for lathe operators in the first stage of analysis (OR 2.36, 95% CI 1.06–5.18), that is attenuated, but still statistically significant in the second stage.

### Analyses on occupational exposure to carcinogens

Analyses for having ever been exposed to one of the 29 selected agents (Table 3) showed significantly increased risks for CD (OR 1.27, 95% CI 1.03–1.57), and borderline significant associations for BAP (OR 1.19, 95% CI 0.96–1.47), CR (OR 1.19, 95% CI 0.94–1.42), PAHs (OR 1.19 95% CI 0.97–1.47), TCE (OR 1.19, 95% CI 0.96–1.46), and TRI (OR 1.21, 95% CI 0.98–1.50).

Considering cumulative exposure, we observed a significantly increased risk of BC for low cumulative exposure to CD (OR 1.34, 95% CI 1.03–1.76), to TCE (OR 1.33, 95% CI 1.02–1.73), and TRI (OR 1.30, 95% CI 1.00–1.73) and a borderline significantly increased risk for low cumulative exposure to BAP (OR 1.26, 95% CI 0.96–1.66), to PB (OR 1.25, 95% CI 0.98–1.61), to PAHs (OR 1.21, 95% CI 0.93–1.59), and to WELD (OR 1.21, 95% CI 0.94–1.56). Moreover, a borderline significantly increased risk was found for high exposure to CD (OR 1.20, 95% CI 0.92–1.56), to CR (OR 1.20, 95% CI 0.94–1.54), and to DET (OR 1.19, 95% CI 0.94–1.52).

Analyses conducted separately by tumour grade showed an involvement of exposure to carcinogens only for low-grade BC (Table 4), while no significant result was found for high-grade BC (Supplementary Table 5). If we consider a dichotomous exposure (ever/never been exposed), for low-grade BC, a significantly increased risk was observed

**Table 2** Significant or borderline significant results of the first stage of the hierarchical model in the job exposure analyses

| ISCO-68 | Label                                     | Cases/controls | OR ML (95% CI)      |
|---------|---|----------------|---------------------|
| 0361    | Chemical engineering technician (general) | 8/1            | 12.65 (1.30–107.12) |
| 3703    | Postmen                                   | 9/2            | 5.48 (1.12–26.83)   |
| 8322    | Tool and die maker                        | 7/1            | 6.03 (1.19–30.63)   |
| 8342    | Lathe operator                            | 50/34          | 1.77 (1.09–2.89)    |
| 9019    | Other rubber and plastics products makers | 7/0            | NA                  |
| 9393    | Spray painter (except construction)       | 9/0            | NA                  |

**Table 3** Results of the multivariable logistic regression models for risk of BC due to exposure (never/ever) and cumulative exposure (no/low/high) to carcinogens

| Cancerogen  | Exposure | Cases/controls | OR (95% CI)      | Exposure | Cases/controls | OR (95% CI)       |
|---|----------|----------------|------------------|----------|----------------|-------------------|
| Aliphatic and alicyclic hydrocarbon solvents (ALHC) | Ever     | 116/119        | 0.94 (0.69–1.27) | High     | 66/60          | 0.99 (0.66–1.49)  |
|   | Never    | 777/859        | Ref.             | Low      | 50/59          | 0.89 (0.58–1.35)  |
|   |          |                |                  | No       | 777/859        | Ref.              |
| Aromatic amines (AA)                                | Ever     | 59/61          | 1.11 (0.75–1.84) | High     | NA             | NA                |
|   | Never    | 828/923        | Ref.             | Low      | NA             | NA                |
|   |          |                |                  | No       | NA             | Ref.              |
| Aromatic hydrocarbon solvents (ARHC)                | Ever     | 212/220        | 0.93 (0.73–1.18) | High     | 124/116        | 0.97 (0.71–1.32)  |
|   | Never    | 681/758        | Ref.             | Low      | 88/104         | 0.86 (0.62–1.19)  |
|   |          |                |                  | No       | 681/758        | Ref.              |
| Arsenic (AS)  | Ever     | 79/81          | 0.99 (0.69–1.41) | High     | 39/41          | 0.85 (0.52–1.41)  |
|   | Never    | 814/897        | Ref.             | Low      | 40/40          | 1.09 (0.67–1.75)  |
|   |          |                |                  | No       | 814/897        | Ref.              |
| Benzene (BENZ)                                      | Ever     | 179/187        | 0.99 (0.77–1.29) | High     | 91/94          | 0.94 (0.67–1.32)  |
|   | Never    | 714/791        | Ref.             | Low      | 88/93          | 1.08 (0.77–1.52)  |
|   |          |                |                  | No       | 714/791        | Ref.              |
| Benzo(a)pyrene (BAP)                                | Ever     | 309/294        | 1.19 (0.96–1.47) | High     | 152/149        | 1.11 (0.84–1.46)  |
|   | Never    | 584/684        | Ref.             | Low      | 157/145        | 1.26 (0.96–1.66)  |
|   |          |                |                  | No       | 584/684        | Ref.              |
| Cadmium (CD)  | Ever     | 359/334        | 1.27 (1.03–1.57) | High     | 183/172        | 1.20 (0.92–1.56)  |
|   | Never    | 534/644        | Ref.             | Low      | 176/162        | 1.34 (1.03–1.76)* |
|   |          |                |                  | No       | 534/644        | Ref.              |
| Chromium (CR)                                       | Ever     | 430/430        | 1.16 (0.94–1.42) | High     | 226/216        | 1.20 (0.94–1.54)  |
|   | Never    | 463/548        | Ref.             | Low      | 204/214        | 1.12 (0.87–1.44)  |
|   |          |                |                  | No       | 463/548        | Ref.              |
| Detergents (DET)                                    | Ever     | 491/508        | 1.16 (0.95–1.43) | High     | 255/258        | 1.19 (0.94–1.52)  |
|   | Never    | 402/470        | Ref.             | Low      | 236/250        | 1.11 (0.87–1.42)  |
|   |          |                |                  | No       | 402/470        | Ref.              |
| Diesel engine exhaust (DEEX)                        | Ever     | 216/256        | 0.81 (0.64–1.02) | High     | 125/129        | 0.92 (0.69–1.24)  |
|   | Never    | 677/722        | Ref.             | Low      | 91/127         | 0.72 (0.52–0.99)  |
|   |          |                |                  | No       | 677/722        | Ref.              |
| Fungicides (FUNG)                                   | Ever     | 82/89          | 0.96 (0.67–1.36) | High     | 47/45          | 1.11 (0.70–1.75)  |
|   | Never    | 811/889        | Ref.             | Low      | 35/44          | 0.77 (0.46–1.28)  |
|   |          |                |                  | No       | 811/889        | Ref.              |
| Gasoline (GASO)                                     | Ever     | 99/116         | 0.9 (0.65–1.23)  | High     | 51/61          | 0.83 (0.54–1.28)  |
|   | Never    | 794/862        | Ref.             | Low      | 48/55          | 1.02 (0.66–1.57)  |
|   |          |                |                  | No       | 794/862        | Ref.              |
| Gasoline engine exhaust (GEEX)                      | Ever     | 226/267        | 0.76 (0.6–0.95)  | High     | 134/146        | 0.81 (0.61–1.07)  |
|   | Never    | 667/711        | Ref.             | Low      | 92/121         | 0.72 (0.52–1)     |
|   |          |                |                  | No       | 667/711        | Ref.              |
| Herbicides (HERB)                                   | Ever     | 77/83          | 0.95 (0.66–1.36) | High     | 42/45          | 0.89 (0.55–1.43)  |
|   | Never    | 816/895        | Ref.             | Low      | 35/38          | 0.97 (0.58–1.63)  |
|   |          |                |                  | No       | 816/895        | Ref.              |
| Insecticides (INSC)                                 | Ever     | 79/86          | 0.95 (0.67–1.36) | High     | 44/44          | 1.06 (0.66–1.7)   |
|   | Never    | 814/892        | Ref.             | Low      | 35/42          | 0.8 (0.48–1.34)   |
|   |          |                |                  | No       | 814/892        | Ref.              |
| Ionizing radiation (IRAD)                           | Ever     | 19/33          | 0.64 (0.33–1.26) | High     | 11/18          | 0.57 (0.24–1.35)  |
|   | Never    | 874/945        | Ref.             | Low      | 8/15           | 0.83 (0.32–2.11)  |
|   |          |                |                  | No       | 874/945        | Ref.              |

**Table 3** (continued)

| Cancerogen                              | Exposure | Cases/controls | OR (95% CI)      | Exposure | Cases/controls | OR (95% CI)      |
|---|----------|----------------|------------------|----------|----------------|------------------|
| Iron (FE)                               | Ever     | 396/399        | 1.11 (0.91–1.37) | High     | 199/207        | 1.07 (0.83–1.38) |
|   | Never    | 497/579        | Ref.             | Low      | 197/192        | 1.17 (0.90–1.51) |
|   |          |                |                  | No       | 497/579        | Ref.             |
| Lead (PB)                               | Ever     | 450/459        | 1.11 (0.91–1.36) | High     | 216/235        | 0.98 (0.77–1.26) |
|   | Never    | 443/519        | Ref.             | Low      | 234/224        | 1.25 (0.98–1.61) |
|   |          |                |                  | No       | 443/519        | Ref.             |
| Low-frequency magnetic fields (MF)      | Ever     | 79/86          | 1.03 (0.84–1.26) | High     | 252/268        | 1.06 (0.83–1.35) |
|   | Never    | 814/892        | Ref.             | Low      | 249/268        | 1.02 (0.80–1.3)  |
|   |          |                |                  | No       | 92/121         | Ref.             |
| Methylene chloride (MCH)                | Ever     | 180/196        | 0.97 (0.76–1.25) | High     | 90/101         | 0.85 (0.61–1.2)  |
|   | Never    | 713/782        | Ref.             | Low      | 90/95          | 1.14 (0.82–1.6)  |
|   |          |                |                  | No       | 713/782        | Ref.             |
| Nickel (NI)                             | Ever     | 403/412        | 1.09 (0.89–1.34) | High     | 210/209        | 1.15 (0.89–1.47) |
|   | Never    | 490/566        | Ref.             | Low      | 193/203        | 1.04 (0.81–1.35) |
|   |          |                |                  | No       | 490/566        | Ref.             |
| Oil mist (OIL)                          | Ever     | 294/324        | 0.98 (0.79–1.22) | High     | 137/162        | 0.92 (0.70–1.22) |
|   | Never    | 599/654        | Ref.             | Low      | 157/162        | 1.07 (0.82–1.4)  |
|   |          |                |                  | No       | 599/654        | Ref.             |
| Other organic solvents (OSOL)           | Ever     | 200/210        | 0.96 (0.75–1.22) | High     | 104/105        | 0.92 (0.67–1.27) |
|   | Never    | 693/768        | Ref.             | Low      | 96/105         | 1.03 (0.74–1.42) |
|   |          |                |                  | No       | 693/768        | Ref.             |
| Perchloroethylene (PER)                 | Ever     | 25/28          | 1.04 (0.56–1.91) | High     | 9/15           | 0.79 (0.31–1.97) |
|   | Never    | 868/950        | Ref.             | Low      | 16/13          | 1.4 (0.62–3.16)  |
|   |          |                |                  | No       | 868/950        | Ref.             |
| Polycyclic aromatic hydrocarbons (PAHs) | Ever     | 344/329        | 1.19 (0.97–1.47) | High     | 175/165        | 1.17 (0.90–1.53) |
|   | Never    | 549/649        | Ref.             | Low      | 169/164        | 1.21 (0.93–1.59) |
|   |          |                |                  | No       | 549/649        | Ref.             |
| Toluene (TOLU)                          | Ever     | 191/210        | 0.92 (0.72–1.18) | High     | 112/107        | 0.99 (0.72–1.37) |
|   | Never    | 344/329        | Ref.             | Low      | 79/103         | 0.87 (0.62–1.23) |
|   |          |                |                  | No       | 702/768        | Ref.             |
| Trichloroethane (TCE)                   | Ever     | 362/358        | 1.18 (0.96–1.46) | High     | 181/185        | 1.08 (0.83–1.41) |
|   | Never    | 531/620        | Ref.             | Low      | 181/173        | 1.33 (1.02–1.73) |
|   |          |                |                  | No       | 531/620        | Ref.             |
| Trichloroethylene (TRI)                 | Ever     | 312/302        | 1.21 (0.98–1.5)  | High     | 157/155        | 1.15 (0.87–1.52) |
|   | Never    | 581/676        | Ref.             | Low      | 155/147        | 1.30 (1.00–1.72) |
|   |          |                |                  | No       | 581/676        | Ref.             |
| Welding fume (WELD)                     | Ever     | 396/399        | 1.11 (0.91–1.37) | High     | 191/202        | 1.03 (0.80–1.33) |
|   | Never    | 497/579        | Ref.             | Low      | 205/197        | 1.21 (0.94–1.56) |
|   |          |                |                  | No       | 479/579        | Ref.             |

Models are adjusted for age, smoking status, intensity of smoking, and study

\*Still statistically significant after FDR correction

for BAP (OR 1.26, 95% CI 1.00–1.61), CD (OR 1.37, 95% CI 1.09–1.73), PAHs (OR 1.29, 95% CI 1.02–1.63), and TRI (OR 1.28, 95% CI 1.01–1.63), while borderline significant results were obtained for CR (OR 1.18, 95% CI 0.94–1.48), DET (OR 1.25, 95% CI 0.99–1.57), FE (OR 1.15, 95% CI 0.92–1.46), PB (OR 1.16, 95% CI 0.92–1.46), and TCE (OR 1.23, 95% CI 0.97–1.55). Considering

cumulative exposure, a significantly increased risk of developing low-grade BC was observed for low exposure to TRI (OR 1.38, 95% CI 1.01–1.87), for high and low exposure to CD (respectively: OR 1.34, 95% CI 1.00–1.79 and OR 1.41, 95% CI 1.04–1.91), while borderline significant results were obtained for low exposure to BAP (OR 1.33, 95% CI 0.98–1.81), DET (OR 1.26, 95% CI



**Table 4** Results of the multivariable logistic regression models for risk of BC due to exposure (never/ever) and cumulative exposure (no/low/high) to carcinogens in low-grade BC

| Carcinogen  | Exposure | Cases/controls | OR (95% CI)       | Exposure | Cases/controls | OR (95% CI)       |
|---|----------|----------------|-------------------|----------|----------------|-------------------|
| Aliphatic and alicyclic hydrocarbon solvents (ALHC) | Ever     | 77/119         | 1.03 (0.74–1.44)  | High     | 42/60          | 1.05 (0.67–1.66)  |
|   | Never    | 479/859        | Ref.              | Low      | 35/59          | 1.01 (0.64–1.6)   |
|   |          |                |                   | No       | 479/859        | Ref.              |
| Aromatic amines (AA)                                | Ever     | 38/61          | 1.18 (0.74–1.82)  | High     | NA             | NA                |
|   | Never    | 515/913        | Ref.              | Low      | NA             | NA                |
|   |          |                |                   | No       | NA             | Ref.              |
| Aromatic hydrocarbon solvents (ARHC)                | Ever     | 126/220        | 0.88 (0.67–1.16)  | High     | 68/116         | 0.87 (0.61–1.25)  |
|   | Never    | 430/758        | Ref.              | Low      | 58/104         | 0.89 (0.62–1.29)  |
|   |          |                |                   | No       | 430/758        | Ref.              |
| Arsenic (AS)  | Ever     | 49/81          | 0.97 (0.65–1.45)  | High     | 22/41          | 0.79 (0.44–1.41)  |
|   | Never    | 507/897        | Ref.              | Low      | 27/40          | 1.17 (0.69–1.98)  |
|   |          |                |                   | No       | 507/897        | Ref.              |
| Benzene (BENZ)                                      | Ever     | 113/187        | 1.05 (0.79–1.4)   | High     | 59/94          | 1.04 (0.71–1.53)  |
|   | Never    | 443/791        | Ref.              | Low      | 54/93          | 1.06 (0.72–1.55)  |
|   |          |                |                   | No       | 443/791        | Ref.              |
| Benzo(a)pyrene (BAP)                                | Ever     | 201/294        | 1.26 (1.00–1.61)  | High     | 98/149         | 1.20 (0.88–1.63)  |
|   | Never    | 355/684        | Ref.              | Low      | 103/145        | 1.33 (0.98–1.81)  |
|   |          |                |                   | No       | 355/684        | Ref.              |
| Cadmium (CD)  | Ever     | 235/334        | 1.37 (1.09–1.73)* | High     | 123/172        | 1.34 (1.00–1.79)  |
|   | Never    | 321/644        | Ref.              | Low      | 112/162        | 1.41 (1.04–1.91)* |
|   |          |                |                   | No       | 321/644        | Ref.              |
| Chromium (CR)                                       | Ever     | 269/430        | 1.18 (0.94–1.48)  | High     | 137/216        | 1.21 (0.91–1.6)   |
|   | Never    | 287/548        | Ref.              | Low      | 132/214        | 1.15 (0.87–1.53)  |
|   |          |                |                   | No       | 287/548        | Ref.              |
| Detergents (DET)                                    | Ever     | 314/508        | 1.25 (0.99–1.57)  | High     | 156/258        | 1.23 (0.93–1.63)  |
|   | Never    | 242/470        | Ref.              | Low      | 158/250        | 1.26 (0.96–1.66)  |
|   |          |                |                   | No       | 242/470        | Ref.              |
| Diesel engine exhaust (DEEX)                        | Ever     | 135/256        | 0.85 (0.65–1.1)   | High     | 74/129         | 0.9 (0.64–1.26)   |
|   | Never    | 421/722        | Ref.              | Low      | 61/127         | 0.79 (0.55–1.13)  |
|   |          |                |                   | No       | 421/722        | Ref.              |
| Fungicides (FUNG)                                   | Ever     | 53/89          | 1.04 (0.7–1.54)   | High     | 33/45          | 1.35 (0.82–2.22)  |
|   | Never    | 503/889        | Ref.              | Low      | 20/44          | 0.72 (0.39–1.33)  |
|   |          |                |                   | No       | 503/889        | Ref.              |
| Gasoline (GASO)                                     | Ever     | 59/116         | 0.89 (0.62–1.28)  | High     | 28/61          | 0.77 (0.46–1.28)  |
|   | Never    | 497/862        | Ref.              | Low      | 31/55          | 1.02 (0.63–1.66)  |
|   |          |                |                   | No       | 497/862        | Ref.              |
| Gasoline engine exhaust (GEEX)                      | Ever     | 135/267        | 0.75 (0.57–0.97)  | High     | 80/146         | 0.77 (0.55–1.06)  |
|   | Never    | 421/711        | Ref.              | Low      | 55/121         | 0.72 (0.49–1.05)  |
|   |          |                |                   | No       | 421/711        | Ref.              |
| Herbicides (HERB)                                   | Ever     | 49/83          | 1.01 (0.67–1.51)  | High     | 28/45          | 1.05 (0.62–1.77)  |
|   | Never    | 507/895        | Ref.              | Low      | 21/38          | 0.95 (0.52–1.74)  |
|   |          |                |                   | No       | 507/895        | Ref.              |
| Insecticides (INSC)                                 | Ever     | 51/86          | 1.03 (0.69–1.54)  | High     | 31/44          | 1.29 (0.77–2.15)  |
|   | Never    | 505/892        | Ref.              | Low      | 20/42          | 0.76 (0.41–1.4)   |
|   |          |                |                   | No       | 505/892        | Ref.              |
| Ionizing radiation (IRAD)                           | Ever     | 13/33          | 0.71 (0.35–1.47)  | High     | 5/18           | 0.32 (0.09–1.10)  |
|   | Never    | 543/945        | Ref.              | Low      | 8/15           | 1.34 (0.53–3.41)  |
|   |          |                |                   | No       | 543/945        | Ref.              |

**Table 4** (continued)

| Carcinogen                              | Exposure | Cases/controls | OR (95% CI)      | Exposure | Cases/controls | OR (95% CI)      |
|---|----------|----------------|------------------|----------|----------------|------------------|
| Iron (FE)                               | Ever     | 249/399        | 1.15 (0.91–1.45) | High     | 126/207        | 1.13 (0.85–1.50) |
|   | Never    | 307/579        | Ref.             | Low      | 123/192        | 1.17 (0.88–1.57) |
|   |          |                |                  | No       | 307/579        | Ref.             |
| Lead (PB)                               | Ever     | 285/459        | 1.16 (0.92–1.46) | High     | 138/235        | 1.06 (0.80–1.41) |
|   | Never    | 271/519        | Ref.             | Low      | 147/224        | 1.28 (0.96–1.69) |
|   |          |                |                  | No       | 271/519        | Ref.             |
| Low-frequency magnetic fields (MF)      | Ever     | 314/536        | 1.07 (0.85–1.35) | High     | 159/268        | 1.11 (0.84–1.46) |
|   | Never    | 242/442        | Ref.             | Low      | 155/268        | 1.04 (0.79–1.37) |
|   |          |                |                  | No       | 242/442        | Ref.             |
| Methylene chloride (MCH)                | Ever     | 115/196        | 1.04 (0.78–1.37) | High     | 56/101         | 0.91 (0.62–1.33) |
|   | Never    | 441/782        | Ref.             | Low      | 59/95          | 1.18 (0.81–1.71) |
|   |          |                |                  | No       | 441/782        | Ref.             |
| Nickel (NI)                             | Ever     | 254/412        | 1.13 (0.9–1.43)  | High     | 126/209        | 1.16 (0.87–1.55) |
|   | Never    | 302/566        | Ref.             | Low      | 128/203        | 1.11 (0.83–1.48) |
|   |          |                |                  | No       | 302/566        | Ref.             |
| Oil mist (OIL)                          | Ever     | 192/324        | 1.06 (0.83–1.35) | High     | 94/162         | 1.04 (0.76–1.43) |
|   | Never    | 364/654        | Ref.             | Low      | 98/162         | 1.08 (0.79–1.46) |
|   |          |                |                  | No       | 364/654        | Ref.             |
| Other organic solvents (OSOL)           | Ever     | 119/210        | 0.93 (0.71–1.23) | High     | 60/105         | 0.88 (0.6–1.28)  |
|   | Never    | 437/768        | Ref.             | Low      | 59/105         | 0.99 (0.69–1.44) |
|   |          |                |                  | No       | 437/768        | Ref.             |
| Perchloroethylene (PER)                 | Ever     | 20/28          | 1.43 (0.76–2.71) | High     | 7/15           | 0.97 (0.36–2.64) |
|   | Never    | 536/950        | Ref.             | Low      | 13/13          | 1.93 (0.82–4.5)  |
|   |          |                |                  | No       | 536/950        | Ref.             |
| Polycyclic aromatic hydrocarbons (PAHs) | Ever     | 225/329        | 1.29 (1.02–1.63) | High     | 117/165        | 1.32 (0.98–1.78) |
|   | Never    | 331/649        | Ref.             | Low      | 108/164        | 1.25 (0.92–1.69) |
|   |          |                |                  | No       | 331/649        | Ref.             |
| Toluene (TOLU)                          | Ever     | 121/210        | 0.97 (0.74–1.28) | High     | 67/107         | 0.97 (0.67–1.4)  |
|   | Never    | 435/768        | Ref.             | Low      | 54/103         | 0.97 (0.67–1.41) |
|   |          |                |                  | No       | 435/768        | Ref.             |
| Trichloroethane (TCE)                   | Ever     | 229/358        | 1.23 (0.97–1.55) | High     | 118/185        | 1.15 (0.86–1.55) |
|   | Never    | 327/620        | Ref.             | Low      | 111/173        | 1.31 (0.97–1.76) |
|   |          |                |                  | No       | 327/620        | Ref.             |
| Trichloroethylene (TRI)                 | Ever     | 199/302        | 1.28 (1.01–1.63) | High     | 99/155         | 1.2 (0.87–1.64)  |
|   | Never    | 357/676        | Ref.             | Low      | 100/147        | 1.38 (1.01–1.87) |
|   |          |                |                  | No       | 357/676        | Ref.             |
| Welding fume (WELD)                     | Ever     | 249/399        | 1.15 (0.91–1.45) | High     | 119/202        | 1.06 (0.79–1.42) |
|   | Never    | 307/579        | Ref.             | Low      | 130/197        | 1.25 (0.94–1.66) |
|   |          |                |                  | No       | 307/579        | Ref.             |

Models are adjusted for age, smoking status, intensity of smoking, and study

\*Still statistically significant after FDR correction

0.96–1.66), to PB (OR 1.28, 95% CI 0.96–1.69), to PAHs (OR 1.25, 95% CI 0.92–1.69), to TCE (OR 1.31, 95% CI 0.97–1.76), and to WELD (OR 1.25, 95% CI 0.94–1.66).

Only the association with CD remained statistically significant after FDR controls.

## Discussion

There is wide agreement in the literature that at least part of BCs share an occupational etiology. This was observed

in several single studies (Kogevinas et al. 2003; Case et al. 1954; Samanic et al. 2008; Purdue et al. 2015; Colt et al. 2014; Lohi et al. 2008) and was confirmed by a recent meta-analysis (Cumberbatch et al. 2017). Most of the occupations that were found to increase the risk for BC in these studies were identified also in several other studies. However, some inconsistencies were present, due in part to the fact that the risk estimates were often computed on small numbers of exposed cases. In fact, several studies pointed out that, when the number of events per variable is less than 10, the regression coefficients are biased (Concato et al. 1995; Peduzzi et al. 1995, 1996), even if, in some cases, this rule could be relaxed (Vittinghoff and McCulloch 2007). Therefore, it is possible that some results reported in the literature are spurious and it appears difficult to identify possible false-positive findings among them.

In our study, we observed an excess risk of BC in chemical engineering technicians, postmen, lathe operators, tool and die makers, as well as an excess BC risk (even if it could not be estimated, due to absence of exposed controls) among rubber and plastic products makers and spray painters. As it was pointed out in the previous studies (Case et al. 1954; Boffetta and Silverman 2001; Vineis and Di Prima 1983; IARC 2010), workers in these occupations were potentially exposed to aromatic amines (mainly: benzidine, 4-aminobiphenyl, b-naphthylamine, and 4-chloro-*o*-toluidine) or to other substances, such as polycyclic aromatic hydrocarbons, diesel engine exhausts, cutting oils, chlorinated hydrocarbons, or other solvents, and it could be speculated that these exposures were causal agents in BC development. In particular, spray painters may have been exposed in the past to spray paints containing water-soluble azo dyes, which have been demonstrated to be metabolized in humans through azoreduction into carcinogenic aromatic amines (Golka et al. 2004a).

To provide a quantitative basis to these associations, we performed a control for false-positive results and an improvement on the estimates within a Bayesian framework, assessing the probability of exposure to carcinogens for each worker through a validated JEM (Kauppinen et al. 2009). After this adjustment, an excess risk of BC could be still observed only for lathe operators, supporting the hypothesis that this occupation, or more generally the work of machine-tool operator, may be an occupation encompassing a high risk for BC. Many studies observed an increased risk of BC for metal workers, and the causal mechanism seems to be related to their exposure to mineral oils that are used to cut, lubricate, and refresh the metal parts wrought in the processes (Colt et al. 2014; Figueroa et al. 2015). Mineral oils, in fact, contain both PAHs and nitrosamines, which are known as human carcinogens (IARC 1987, 2000, 2010).

The second aim of our study was to focus the attention on carcinogens rather than on occupations. Therefore, we employed a validated JEM to assess the association of BC with having ever been exposed and with cumulative exposure to selected carcinogens, using information on subjects' jobs.

The strongest result obtained was a 37% increase in BC risk due to CD exposure (41% for low cumulative exposure). CD was shown to be carcinogenic in humans through four different main mechanisms: aberrant gene expression, inhibition of DNA damage repair, inhibition of apoptosis, and induction of oxidative stress (Joseph 2009; Waisberg et al. 2003); CD has been classified as carcinogenic to humans (group 1) by the IARC, which stated that there is sufficient evidence for a causal association with lung cancer, and limited evidence for prostate cancer and kidney cancer (IARC 2012). Only a few *in vivo* models (Waalkes et al. 2000; Talaska et al. 1990) have been developed to assess the carcinogenicity of CD for BC, with contrasting results. Moreover, in an ecological study performed in Belgium to identify a possible relationship between cancer incidence and cadmium emissions from zinc smelters, no association was found with bladder (and any other) cancer (Verhoeven et al. 2011). Nevertheless, Feki-Tounsi and Hamza-Caffai (Feki-Tounsi and Hamza-Chaffai 2014) published a review in which they concluded that results from *in vitro* studies (Sens et al. 2004; Larson et al. 2010) and from previous epidemiological studies (Feki-Tounsi et al. 2013; Kellen et al. 2007) would suggest an association between CD exposure and BC development.

We also found an excess risk of BC in subjects with low cumulative exposure to PB and TRI, when compared to non-exposed subjects. In a Finnish study conducted by Lohi et al. (2008), exposure to Chlorinated Aliphatic Hydrocarbons (CHC), mainly TRI e TCE, was associated with an increased risk of BC, in particular for female laboratory assistants, chemical industry laborers and shoemakers, and for male transport operators like taxi and bus drivers. Moreover, many animal experiments confirmed TRI and PER to be carcinogenic agents for urological cancers (Golka et al. 2004b). Our results were less strong than those observed in the previous studies, possibly because the increase in risk awareness and the implementation of control measures have reduced considerably the level of exposure to solvents in the last decades [for example, exposure to tetrachlorethylene in dry cleaning has decreased from 350 to 700 mg/m<sup>3</sup> in the '70s to 70–350 mg/m<sup>3</sup> in '80s (IARC 1995)].

Regarding the association with PB exposure, a recent review from Assi et al. (2016) on the effects of PB on human and animal health showed that PB is a possible cause of urinary cancer development (in particular for the kidney).

We also analyzed the risk of BC due to carcinogenic exposure by tumour grade. We observed that the carcinogens

considered had a higher effect on the risk of low-grade BC, while it seems that they had only a marginal role in the risk of high-grade BC. In fact, together with the already observed increased risk due to CD, TRI, TCE, PAHs, and PB exposure, a statistically or borderline-statistically significant increased risk of developing low-grade BC was found also for BAP, CR, and DET exposure, while no association was found for high-grade BC. As a partial rationale for this, a recent paper (Van Batavia et al. 2014) showed that low-grade BC arises mainly from intermediate cells, while Keratin5-expressing basal cells are likely progenitors of high-grade BC, more prone to undergo a *TP53* mutation. Therefore, it is not surprising that carcinogens have different effects on these two different types of cells. To our knowledge, no other studies investigated the effect of carcinogens on BC risk by tumour grade, except for an old study which considered the association between BC and having worked in occupations bearing a high a priori risk for this cancer, and did not find any difference in the association with respect to grade (about 50% risk excess in both categories) (Sturgeon et al. 1994). Therefore, new studies are needed to further investigate this topic.

The main limitation of the present study lies in the fact that it is based on a hospital case–control design; indeed, it could be difficult to judge how much controls reflected the distribution of the population that generated cases. For this reason, we tested, if a socio-economic difference between cases and controls was present evaluating the frequency distribution of ISCO-68 macro-classes in cases and controls and, since there was no difference, it is unlikely that the observed associations were due to a possible higher socio-economic status of controls. Moreover, the use of hospital controls could bring to a not adequate representation of the underlying study base that generated the cases, although the percentage of Italian employees by broad occupational groups in those years is similar to that presented in the controls (<http://dati.istat.it>).

Another limitation is the use of a JEM to assign occupational exposure to the study population, which implies higher non-differential misclassification of the exposure than more detailed information at the individual level, with the consequence of a risk attenuation. However, the use of a JEM can also be a strength, given that it reduces the possibility of differential misclassification due to recall bias, that is more severe, due to its differential nature, and may lead to under- or overestimation of the true risk. Recall bias was also reduced keeping into account only jobs lasted for at least 2 years during the lifetime.

The construction of the cumulative dose of exposure through intensity weighted by probability may have also led to an imperfect classification of the subjects in low- or high cumulative exposure, and in particular to an underestimation of the true cumulative exposure for subjects with low

probability and high intensity. This fact could be partially explained why some associations were stronger in subjects with lower exposure than in highly exposed subjects.

A strength of the study is the multi-centre approach, that has increased not only the sample size, but also the representativeness of the results, describing two different industrial and then post-industrial environments. Conversely, the two studies had not exactly the same design and this could lead to some heterogeneity.

Moreover, with respect to the previous studies, we performed an improvement of the plausibility and robustness of the risk estimates and a rigorous control of type I error, which is always present when performing many independent tests, employing semi-Bayesian analyses. Finally, we studied both occupation and occupational exposure to carcinogens, to provide a more detailed picture of BC risk among workers.

In conclusion, our study showed that several occupations, especially those involving rubber, plastic, and mechanical work, may bear higher risk of developing BC, as well as occupational exposure to several carcinogens, such as cadmium, lead, and chlorinated aliphatic hydrocarbons, which are not yet officially recognized as carcinogens for BC. These findings suggest that monitoring workplaces to prevent exposure to carcinogenic agents is still a very important task, which should be still given adequate importance in public health policies.

**Acknowledgements** We would like to thank Dr. Timo Kauppinen for giving us the opportunity to use the NOCCA JEM. The study was partly funded through an agreement between the Universities of Verona and Brescia, Italy.

## References

- AIOM AIRTUM (2016) I numeri del cancro in Italia 2016. Intermedia Editore, Brescia
- Assi MA, Hezme MN, Haron AW, Sabri MY, Rajion MA (2016) The detrimental effects of lead on human and animal health. *Vet World* 6:660–671
- Babjuk M, Burger M, Zigeuner R, Shariat SF, van Rhijn BW, Compérat E, Sylvester RJ, Kaasinen E, Böhle A, Palou Redorta J, Rouprêt M (2013) European Association of Urology. EAU guidelines on non-muscle-invasive urothelial carcinoma of the bladder: update 2013. *Eur Urol* 64(4):639–653. <https://doi.org/10.1016/j.eururo.2013.06.003>
- Benjamini Y, Hochberg Y (1995) Controlling the false discovery rate: a practical and powerful approach to multiple testing. *J R Stat Soc Ser B (Methodological)* 57(1):289–300
- Boffetta P, Silverman DT (2001) A meta-analysis of bladder cancer and diesel exhaust exposure. *Epidemiology* 12(1):125–130
- Boiano JM, Hull RD (2001) Development of a National Occupational Exposure Survey and database associated with NIOSH hazard surveillance initiatives. *Appl Occup Environ Hyg* 16(2):128–134
- Case R, Hosker ME (1954) Tumour of the urinary bladder as an occupational disease in the rubber industry in England and Wales. *Br J Ind Med* 8(2):39–50

- Case RA, Hosker ME, McDonald DB, Pearson JT (1954) Tumours of the urinary bladder in workmen engaged in the manufacture and use of certain dyestuff intermediates in the British chemical industry. I. The role of aniline, benzidine, alpha-naphthylamine, and beta-naphthylamine. *Br J Int Med* 11:75–104
- Colt JS, Friesen MC, Stewart PA, Donguk P, Johnson A, Schwenn M, Karagas MR, Armenti K, Waddell R, Verrill C, Ward MH, Freeman LE, Moore LE, Koutros S, Baris D, Silverman DT (2014) A case-control study of occupational exposure to metalworking fluids and bladder cancer risk among men. *Occup Environ Med* 71(10):667–674. <https://doi.org/10.1136/oemed-2013-102056>
- Concato J, Peduzzi P, Holford TR, Feinstein AR (1995) Importance of events per independent variable in proportional hazards analysis. I. Background, goals, and general strategy. *J Clin Epidemiol* 48(12):1495–1501
- Corbin M, Richiardi L, Vermeulen R, Kromhout H, Merletti F, Peters S, Simonato L, Steenland K, Pearce N, Maule M (2012) Hierarchical regression for multiple comparisons in a case-control study of occupational risks for lung cancer. *PLoS One* 7(6):1–8. <https://doi.org/10.1371/journal.pone.0038944>
- Cumberbatch MG, Windsor-Sheppard B, Catto JW (2017) The contemporary landscape of occupational bladder cancer within the United Kingdom: a meta-analysis of risks over the last 80 years. *BJU Int* 119(1):100–109. <https://doi.org/10.1111/bju.13561>
- d'Errico A, Pasian S, Baratti A, Zanelli R, Alfonso S, Gilardi L, Beatrice F, Bena A, Costa G (2009) A case-control study on occupational risk factors for sino-nasal cancer. *Occup Environ Med* 66(7):448–455. <https://doi.org/10.1136/oem.2008.041277>
- Falcone U, Gilardi L, Santoro S, Oreggia M, Marighella M, Coffano ME (2013) MATline, a job-exposure matrix for the prevision of exposure to carcinogens: new functions and potential applications. *Epidemiol Prev* 37(1):60–66
- Feki-Tounsi M, Hamza-Chaffai A (2014) Cadmium as a possible cause of bladder cancer: a review of accumulated evidence. *Environ Sci Pollut Res Int* 21(18):10561–10573. <https://doi.org/10.1007/s11356-014-2970-0>
- Feki-Tounsi M, Olmedo P, Gil F, Khelifi R, Mhiri MN, Rebai A, HamzaChaffai A (2013) Cadmium in blood of Tunisian men and risk of bladder cancer: interactions with arsenic exposure and smoking. *Environ Sci Pollut Res Int* 20(10):7204–7213. <https://doi.org/10.1007/s11356-013-1716-8>
- Figueroa JD, Koutros S, Colt JS, Kogevinas M, Garcia-Closas M, Real FX, Friesen MC, Baris D, Stewart P, Schwenn M, Johnson A, Karagas MR, Armenti KR, Moore LE, Schned A, Lenz P, Prokunina-Olsson L, Banday AR, Paquin A, Ylaya K, Chung JY, Hewitt SM, Nickerson ML, Tardón A, Serra C, Carrato A, García-Closas R, Lloreta J, Malats N, Fraumeni JF Jr, Chanock SJ, Chatterjee N, Rothman N, Silverman DT (2015) Modification of occupational exposures on bladder cancer risk by common genetic polymorphisms. *J Natl Cancer Inst* 107(11):djv223
- Golka K, Kopps S, Myslak ZW (2004a) Carcinogenicity of azo colorants: influence of solubility and bioavailability. *Toxicol Lett* 151(1):203–210
- Golka K, Wiese A, Assennato G, Bolt HM (2004b) Occupational exposure and urological cancer. *World J Urol* 21(6):382–391
- Greenland S (1994) Hierarchical regression for epidemiologic analysis of multiple exposures. *Environ Health Perspect* 102(suppl 8):33–39
- Hrudey SE, Backer L, Humpage AR, Krasner SW, Michaud DS, Moore LE, Singer PC, Stanford BD (2015) Evaluating evidence for association of human bladder cancer with drinking water chlorination disinfection by-products. *J Toxicol Environ Health* 18(5):213–241. <https://doi.org/10.1080/10937404.2015.1067661>
- IARC (1987) Polynuclear aromatic compounds, part 2: carbon blacks, mineral oil (lubricant base oils and derived products) and some nitroarenes. IARC, Lyon
- IARC Monographs on the Evaluation of Carcinogenic Risks to Humans (1995) Volume 63: dry cleaning, some chlorinated solvents and other industrial chemicals. IARC, Lyon
- IARC Monographs on the Evaluation of Carcinogenic Risks to Humans (2000) Volume 77: some industrial chemicals. IARC, Lyon
- IARC Monographs on the Evaluation of Carcinogenic Risks to Humans (2010) Volume 98: painting, firefighting, and shiftwork. IARC, Lyon
- IARC Monographs on the Evaluation of Carcinogenic Risks to Humans (2012) Volume 100F: chemical agents and related occupations. IARC, Lyon
- International Labour Office (1968) ISCO. International Standard Classification of Occupations. International Labor Organisation, Geneva
- ITACAN (2018) Tumori in Italia, Versione 2.0. Associazione Italiana dei Registri TUMori <http://www.registri-tumori.it>. Accessed Jan 2018
- Joseph P (2009) Mechanisms of cadmium carcinogenesis. *Toxicol Appl Pharmacol* 238(3):272–279. <https://doi.org/10.1016/j.taap.2009.01.011>
- Kauppinen T, Heikkilä P, Plato N, Woldbaek T, Lenvik H, Hansen J, Kristjansson V, Pukkala E (2009) Construction of job-exposure matrices for the Nordic Occupational Cancer Study (NOCCA). *Acta Oncol* 48(5):791–800. <https://doi.org/10.1080/02841860902718747>
- Kellen E, Zeegers MP, Hond ED, Buntinx F (2007) Blood cadmium may be associated with bladder carcinogenesis: the Belgian case-control study on bladder cancer. *Cancer Detect Prev* 31(1):77–82
- Kogevinas M, Kauppinen T, Boffetta P, Saracci R (1998) Estimation of the burden of occupational cancer in Europe. Final report to the European commission of a project funded by the programme in Europe Against Cancer. Tech rep, Institut Municipal d'Investigacio Medica
- Kogevinas M, t'Mannetje A, Cordier S, Ranft U, González CA, Vineis P, Chang-Claude J, Lynge E, Wahrendorf J, Tzonou A, Jöckel KH, Serra C, Porru S, Hours M, Greiser E, Boffetta P (2003) Occupation and bladder among men in Western Europe. *Cancer Causes Control* 14(10):907–914
- Larson J, Yasmin T, Sens DA, Zhou XD, Sens MA, Garrett SH, Dunlevy JR, Cao L, Somji S (2010) SPARC gene expression is repressed in human urothelial cells (UROtsa) exposed to or malignantly transformed by cadmium or arsenite. *Toxicol Lett* 199(2):166–172
- Lohi J, Kyyronen P, Kauppinen T, Kujala V, Pukkala E (2008) Occupational exposure to solvents and gasoline and risk of cancers in the urinary tract among finnish workers. *Am J Ind Med* 51(9):668–672. <https://doi.org/10.1002/ajim.20606>
- McGeoghegan D, Binks K (2000) The mortality and cancer morbidity experience of workers at the Capenhurst uranium enrichment facility 1946–95. *J Radiol Prot* 20(4):381–401
- Negri E, La Vecchia C (2001) Epidemiology and prevention of bladder cancer. *Eur J Cancer Prev* 10:7–14
- Peduzzi P, Concato J, Feinstein AR, Holford TR (1995) Importance of events per independent variable in proportional hazards regression analysis. II. Accuracy and precision of regression estimates. *J Clin Epidemiol* 48(12):1503–1510
- Peduzzi P, Concato J, Kemper E, Holford TR, Feinstein AR (1996) A simulation study of the number of events per variable in logistic regression analysis. *J Clin Epidemiol* 49(12):1373–1379
- Porru S, Aulenti V, Donato F, Boffetta P, Fazioli R, Cosciani Cunico S, Alessio L (1996) Bladder cancer and occupation: a case-control study in Northern Italy. *Occup Environ Med* 53(1):6–10
- Porru S, Pavanello S, Carta A, Arici C, Simeone C, Izzotti A, Mastrofrancesco G (2014) Complex relationships between occupation, environment, DNA adducts, genetic polymorphisms and bladder

- cancer in a case–control study using a structural equation modeling. *PLoS One* 9(4):e94566
- Purdue MP, Hutchings SJ, Rushton L, Silverman DT (2015) The proportion of cancer attributable to occupational exposures. *Ann Epidemiol* 25(3):188–192. <https://doi.org/10.1016/j.annepidem.2014.11.009>
- Ricceri F, Guarrera S, Sacerdote C, Polidoro S, Allione A, Fontana D, Destefanis P, Tizzani A, Casetta G, Cucchiareale G, Vineis P, Matullo G (2010) ERCC1 haplotypes modify bladder cancer risk: a case–control study. *DNA Repair* 9(2):191–200. <https://doi.org/10.1016/j.dnarep.2009.12.002>
- Rota M, Bosetti C, Boccia S, Boffetta P, La Vecchia C (2014) Occupational exposures to polycyclic aromatic hydrocarbons and respiratory and urinary tract cancers: an updated systematic review and a meta-analysis to 2014. *Arch Toxicol* 88(8):1479–1490. <https://doi.org/10.1007/s00204-014-1296-5>
- Rushton L, Hutchings S, Brown T (2008) The burden of cancer at work: estimation as the first step to prevention. *Occup Environ Med* 65(12):789–800
- Sacerdote C, Matullo G, Polidoro S, Gamberini S, Piazza A, Karagas MR, Rolle L, De Stefani P, Casetta G, Morabito F, Vineis P, Guarrera S (2007) Intake of fruits and vegetables and polymorphisms in DNA repair genes in bladder cancer. *Mutagenesis* 22(4):281–285
- Sacerdote C, Guarrera S, Ricceri F, Pardini B, Polidoro S, Allione A, Critelli R, Russo A, Andrew AS, Ye Y, Wu X, Kiemeny LA, Bosio A, Casetta G, Cucchiareale G, Destefanis P, Gontero P, Rolle L, Zitella A, Fontana D, Vineis P, Matullo G (2013) Polymorphisms in the *XRCC1* gene modify survival of bladder cancer patients treated with chemotherapy. *Int J Cancer* 133(8):2004–2009
- Saint-Jacques N, Parker L, Brown P, Dummer T (2014) Arsenic in drinking water and urinary tract cancers: a systematic review of 30 years of epidemiological evidence. *Environ Health* 13:44. <https://doi.org/10.1186/1476-069X-13-44>
- Samanic CM, Kogevinas M, Silverman DT, Tardón A, Serra C, Malats N, Real FX, Carrato A, García-Closas R, Sala M, Lloreta J, Rothman N, Dosemeci M (2008) Occupation and bladder cancer in a hospital-based case–control study in Spain. *Occup Environ Med* 65(5):347–353
- Schottenfeld D, Fraumeni JF (2006) *Cancer epidemiology and prevention*, 3rd edn. Oxford University press, Oxford
- Sens DA, Park S, Gurel V, Sens MA, Garrett SH, Somji S (2004) Inorganic cadmium- and arsenite-induced malignant transformation of human bladder urothelial cells. *Toxicol Sci* 79(1):56–63
- Stewart BW, Wild CP (2014) *World cancer report 2014*. IARC, Lyon
- Sturgeon SR1, Hartge P, Silverman DT, Kantor AF, Linehan WM, Lynch C, Hoover RN (1994) Associations between bladder cancer risk factors and tumor stage and grade at diagnosis. *Epidemiology* 5(2):218–225
- Talaska G, Dooley KL, Kadlubar FF (1990) Detection and characterization of carcinogen-DNA adducts in exfoliated urothelial cells from 4-aminobiphenyl-treated dogs by <sup>32</sup>P-postlabelling and subsequent thin-layer and high-pressure liquid chromatography. *Carcinogenesis* 11(4):639–646
- Van Batavia J, Yamany T, Molotkov A, Dan H, Mansukhani M, Batourina E, Schneider K, Oyon D, Dunlop M, Wu XR, Cordon-Cardo C, Mendelsohn C (2014) Bladder cancers arise from distinct urothelial sub-populations. *Nat Cell Biol* 16(10):982–991. <https://doi.org/10.1038/ncb3038>
- Van Osch FHM, Jochems SHJ, Van Schooten FJ, Bryan RT, Zeegers MP (2016) Quantified relations between exposure to tobacco smoking and bladder cancer risk: a meta-analysis of 89 observational studies. *Int J Epidemiol* 45(3):857–870. <https://doi.org/10.1093/ije/dyw044>
- Verhoeven RH, Louwman MW, Buntinx F, Botterweck AM, Lousbergh D, Faes C, Coebergh JW (2011) Variation in cancer incidence in northeastern Belgium and southeastern Netherlands seems unrelated to cadmium emission of zinc smelters. *Eur J Cancer Prev* 20(6):549–555
- Vineis P, Di Prima S (1983) Cutting oil and bladder cancer. *Scand J Work Environ Health* 9(5):449–450
- Vineis P, Pirastu R (1997) Aromatic amines and cancer. *Cancer Causes Control* 8(3):346–355
- Vittinghoff E, McCulloch CE (2007) Relaxing the rule of ten events per variable in logistic and Cox regression. *Am J Epidemiol* 165(6):710–718
- Waalkes MP, Rehm S, Cherian MG (2000) Repeated cadmium exposures enhance the malignant progression of ensuing tumors in rats. *Toxicol Sci* 54(1):110–120
- Waisberg M, Joseph P, Hale B, Beyersmann D (2003) Molecular and cellular mechanisms of cadmium carcinogenesis. *Toxicology* 192:95–117
- Weiss HA, Darby SC, Doll R (1994) Cancer mortality following X-ray treatment for ankylosing spondylitis. *Int J Cancer* 59(3):327–338
- Witte JS, Greenland S, Haile RW, Bird CL (1994) Hierarchical regression analysis applied to a study of multiple dietary exposures and breast cancer. *Epidemiology* 5(6):612–621

**Publisher's Note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

**Supplementary Table 1. Agents identified by the International Agency for Research on Cancer (IARC) as possibly involved in the carcinogenetic process of BC or other cancers of the genitourinary system**

*GROUP 1 – Certainly carcinogenic to humans*

Arsenic (AS)

Aromatic amines (AA: 4-Aminobiphenyl. benzidine. 2-Naphthylamine. Ortho-Toluidine)

Benzo[a]pyrene (BAP)

Chlorinated hydrocarbon solvents (CHC: Trichloroethylene. 1.2-Dichloropropane).

Welding fumes (WELD)

*GROUP 2 – Probably or possibly carcinogenic to humans*

Aliphatic and alicyclic hydrocarbon solvents (ALHC)

Aromatic hydrocarbon solvents (ARHC)

Benzene (BENZ)

Cadmium (CD)

Chromium (CR)

Detergents (DET)

Diesel engine exhausts (DEEX)

Fungicides (FUNG)

Gasoline (GASO)

Gasoline engine exhaust (GEEX)

Herbicides (HERB)

Insecticides (INSC)

Ionizing radiation (IRAD)

Iron (FE)

Lead (PB)

Low frequency magnetic fields (MF)

Methylene chloride (MCH)

Nickel (NI)

Oil mist (OIL)

Other organic solvents (OSOL)

Perchloroethylene (PER)

Toluene (TOLU)

Trichloroethane (TCE)

Trichloroethylene (TRI)

**Supplementary Table 2. Years of exposure weighted by probability and intensity equivalent to 1 year of cumulative exposure.**

|                    | <b>Intensity</b>        |               |             | <b>Years of exposure</b> |     |     |
|--------------------|-------------------------|---------------|-------------|--------------------------|-----|-----|
|                    | <b>Low</b>              | <b>Medium</b> | <b>High</b> |                          |     |     |
| <b>Probability</b> | <b>Assigned weights</b> |               |             |                          |     |     |
| Low                | 0.3                     | 0.3           | 0.3         | 3.5                      | 3.5 | 3.5 |
| Medium             | 0.5                     | 0.7           | 0.7         | 2.0                      | 1.5 | 1.5 |
| High               | 0.7                     | 1.0           | 1.0         | 1.5                      | 1.0 | 1.0 |



**Supplementary Table 3. Results for the risk of BC due to occupations performed for at least 2 years with at least 5 cases. Models are adjusted for age, smoking status, intensity of smoking and study.**

| NCSO | Label   | Cases/<br>controls | ML<br>OR [95% CI]   | HR $\tau=0.76$   |                  | HR $\tau=0.59$   |                  | HR $\tau=0.41$ |          | HR $\tau=0.23$ |          |
|------|---|--------------------|---------------------|------------------|------------------|------------------|------------------|----------------|----------|----------------|----------|
|      |   |                    |                     | OR               | [95% CI]         | OR               | [95% CI]         | OR             | [95% CI] | OR             | [95% CI] |
| 3322 | Mechanical draughtsman                        | 18/14              | 1.85 [0.85-4.01]    | 1.34 [0.92-1.95] | 1.33 [0.92-1.92] | 1.3 [0.92-1.84]  | 1.22 [0.9-1.63]  |                |          |                |          |
| 3351 | Mechanical Engineering Technician             | 5/6                | 0.94 [0.27-3.2]     | 0.98 [0.56-1.73] | 0.99 [0.57-1.7]  | 1 [0.61-1.63]    | 1.02 [0.7-1.47]  |                |          |                |          |
| 3361 | Chemical Engineering Technician               | 8/1                | 12.65 [1.30-107.12] | 2.43 [0.65-6.32] | 1.71 [0.58-3.32] | 1.32 [0.47-2.39] | 1.05 [0.66-1.65] |                |          |                |          |
| 3399 | Other Engineering Technicians.                | 7/8                | 1.05 [0.36-3.11]    | 1.03 [0.62-1.72] | 1.03 [0.63-1.69] | 1.04 [0.66-1.63] | 1.04 [0.73-1.47] |                |          |                |          |
| 3721 | Auxiliary Nurse.                              | 6/8                | 0.89 [0.26-3.04]    | 0.96 [0.54-1.7]  | 0.97 [0.56-1.67] | 0.98 [0.6-1.61]  | 1.01 [0.7-1.45]  |                |          |                |          |
| 3842 | Computer programmer                           | 5/6                | 0.9 [0.25-3.23]     | 0.96 [0.54-1.74] | 0.97 [0.56-1.71] | 0.99 [0.6-1.63]  | 1.01 [0.7-1.47]  |                |          |                |          |
| 3328 | Technical education teacher (second level)    | 6/2                | 4.16 [0.73-23.8]    | 1.73 [0.81-3.68] | 1.61 [0.8-3.24]  | 1.43 [0.79-2.59] | 1.2 [0.8-1.8]    |                |          |                |          |
| 1329 | Other secondary education teachers            | 5/4                | 2.29 [0.55-9.54]    | 1.42 [0.75-2.7]  | 1.38 [0.75-2.53] | 1.29 [0.76-2.21] | 1.16 [0.79-1.71] |                |          |                |          |
| 1332 | First-level education teacher                 | 6/3                | 1.83 [0.43-7.83]    | 1.29 [0.67-2.49] | 1.27 [0.68-2.34] | 1.21 [0.71-2.08] | 1.12 [0.76-1.65] |                |          |                |          |
| 2111 | General managers                              | 28/27              | 1.2 [0.67-2.15]     | 1.09 [0.82-1.46] | 1.09 [0.82-1.45] | 1.09 [0.83-1.44] | 1.08 [0.84-1.38] |                |          |                |          |
| 2121 | Production Manager (except farm)              | 9/3                | 1.72 [0.48-6.22]    | 1.27 [0.7-2.3]   | 1.25 [0.71-2.19] | 1.21 [0.73-2]    | 1.13 [0.78-1.63] |                |          |                |          |
| 2199 | Other managers                                | 14/11              | 1.11 [0.45-2.72]    | 1.06 [0.69-1.62] | 1.06 [0.69-1.6]  | 1.05 [0.71-1.56] | 1.05 [0.76-1.44] |                |          |                |          |
| 2101 | Government Executive Official                 | 8/7                | 0.73 [0.27-2.01]    | 0.88 [0.54-1.41] | 0.89 [0.56-1.41] | 0.91 [0.59-1.4]  | 0.96 [0.68-1.34] |                |          |                |          |
| 2311 | Bookkeeper                                    | 6/4                | 1.18 [0.28-4.93]    | 1.08 [0.57-2.07] | 1.08 [0.59-1.99] | 1.07 [0.63-1.84] | 1.06 [0.72-1.55] |                |          |                |          |
| 2394 | Finance clerk                                 | 5/15               | 0.45 [0.15-1.33]    | 0.71 [0.42-1.18] | 0.73 [0.44-1.19] | 0.77 [0.49-1.22] | 0.87 [0.61-1.24] |                |          |                |          |
| 2604 | Bus Conductor.                                | 6/3                | 2.67 [0.62-11.47]   | 1.45 [0.76-2.79] | 1.38 [0.75-2.55] | 1.25 [0.73-2.14] | 1.07 [0.72-1.58] |                |          |                |          |
| 3703 | Postmen                                       | 9/2                | 5.48 [1.12-26.83]   | 1.97 [0.98-3.97] | 1.82 [0.94-3.51] | 1.58 [0.9-2.79]  | 1.27 [0.85-1.89] |                |          |                |          |
| 2704 | messenger                                     | 15/16              | 1 [0.48-2.1]        | 1 [0.69-1.43]    | 1 [0.7-1.43]     | 0.99 [0.7-1.41]  | 0.98 [0.71-1.35] |                |          |                |          |
| 3914 | Storeroom clerk                               | 8/10               | 0.75 [0.28-2]       | 0.89 [0.56-1.41] | 0.9 [0.57-1.41]  | 0.92 [0.6-1.39]  | 0.96 [0.69-1.34] |                |          |                |          |
| 3922 | Material planning clerk                       | 7/12               | 0.8 [0.3-2.08]      | 0.91 [0.57-1.44] | 0.92 [0.59-1.43] | 0.93 [0.62-1.41] | 0.97 [0.7-1.35]  |                |          |                |          |
| 3923 | Production Planning Clerk.                    | 5/5                | 1.38 [0.38-5.09]    | 1.16 [0.64-2.11] | 1.15 [0.65-2.03] | 1.13 [0.68-1.87] | 1.08 [0.75-1.58] |                |          |                |          |
| 3931 | Office clerk                                  | 58/53              | 1.26 [0.82-1.94]    | 1.12 [0.91-1.39] | 1.12 [0.91-1.38] | 1.12 [0.91-1.38] | 1.11 [0.91-1.35] |                |          |                |          |
| 4102 | Working Proprietor (Wholesale Trade).         | 9/10               | 1.11 [0.43-2.85]    | 1.06 [0.67-1.66] | 1.06 [0.68-1.63] | 1.06 [0.7-1.59]  | 1.05 [0.76-1.46] |                |          |                |          |
| 4103 | Working proprietor (retail trade)             | 22/20              | 0.88 [0.47-1.67]    | 0.95 [0.69-1.29] | 0.95 [0.7-1.29]  | 0.95 [0.71-1.29] | 0.97 [0.75-1.26] |                |          |                |          |
| 4219 | Other sales supervisors                       | 7/5                | 1.64 [0.47-5.78]    | 1.25 [0.7-2.23]  | 1.23 [0.71-2.14] | 1.19 [0.72-1.96] | 1.12 [0.77-1.62] |                |          |                |          |
| 4322 | Commercial traveller                          | 19/32              | 0.59 [0.32-1.08]    | 0.78 [0.58-1.05] | 0.78 [0.58-1.05] | 0.8 [0.6-1.06]   | 0.84 [0.66-1.09] |                |          |                |          |
| 4323 | Manufacturers' agent                          | 17/18              | 0.75 [0.35-1.61]    | 0.88 [0.61-1.27] | 0.89 [0.62-1.27] | 0.9 [0.64-1.27]  | 0.94 [0.7-1.26]  |                |          |                |          |
| 4412 | Insurance salesman                            | 6/6                | 1.1 [0.31-3.83]     | 1.05 [0.59-1.87] | 1.05 [0.61-1.82] | 1.05 [0.64-1.73] | 1.04 [0.72-1.51] |                |          |                |          |
| 4512 | Wholesale trade salesman                      | 7/3                | 2.97 [0.71-12.45]   | 1.58 [0.83-3.01] | 1.51 [0.82-2.78] | 1.39 [0.81-2.38] | 1.2 [0.82-1.77]  |                |          |                |          |
| 4513 | Cook, except private service                  | 26/25              | 1.21 [0.67-2.19]    | 1.1 [0.82-1.47]  | 1.1 [0.82-1.46]  | 1.09 [0.83-1.45] | 1.08 [0.84-1.39] |                |          |                |          |
| 4519 | Other salesmen, shop assistants, demonstrator | 7/7                | 1.22 [0.41-3.62]    | 1.1 [0.66-1.84]  | 1.1 [0.67-1.8]   | 1.09 [0.69-1.71] | 1.07 [0.75-1.52] |                |          |                |          |
| 4522 | Street vendor                                 | 16/18              | 0.83 [0.41-1.68]    | 0.92 [0.65-1.3]  | 0.92 [0.66-1.3]  | 0.93 [0.67-1.29] | 0.96 [0.72-1.27] |                |          |                |          |

| ISCO | Label   | Cases/<br>controls | ML<br>OR | HR $\tau=0.76$ |         | HR $\tau=0.59$ |         | HR $\tau=0.41$ |         | HR $\tau=0.23$ |         |             |
|------|---|--------------------|----------|----------------|---------|----------------|---------|----------------|---------|----------------|---------|-------------|
|      |   |                    |          | OR             | 95% CI] | OR             | 95% CI] | OR             | 95% CI] | OR             | 95% CI] |             |
| 6105 | Working proprietor (café, bar and snack bar)  | 6/7                | 0.88     | [0.29-2.64]    | 0.95    | [0.57-1.6]     | 0.96    | [0.58-1.58]    | 0.97    | [0.62-1.54]    | 1       | [0.7-1.42]  |
| 6312 | Head cook   | 8/3                | 2.65     | [0.53-13.34]   | 1.48    | [0.73-3.02]    | 1.42    | [0.73-2.75]    | 1.31    | [0.74-2.32]    | 1.16    | [0.78-1.73] |
| 6313 | Cook, except private service  | 6/5                | 1.34     | [0.41-4.38]    | 1.14    | [0.66-1.99]    | 1.14    | [0.67-1.93]    | 1.12    | [0.69-1.81]    | 1.08    | [0.75-1.56] |
| 6321 | Waiter, general   | 9/7                | 2.08     | [0.62-6.93]    | 1.38    | [0.79-2.42]    | 1.35    | [0.79-2.31]    | 1.29    | [0.8-2.09]     | 1.17    | [0.81-1.69] |
| 6325 | Bartender   | 18/13              | 1.18     | [0.52-2.64]    | 1.08    | [0.73-1.6]     | 1.08    | [0.74-1.59]    | 1.08    | [0.75-1.55]    | 1.07    | [0.79-1.44] |
| 6522 | Charworker  | 12/5               | 2.08     | [0.75-5.82]    | 1.4     | [0.86-2.27]    | 1.37    | [0.86-2.2]     | 1.32    | [0.85-2.04]    | 1.2     | [0.85-1.69] |
| 6703 | Barber-hairdresser  | 11/16              | 1.04     | [0.44-2.48]    | 1.03    | [0.68-1.56]    | 1.03    | [0.68-1.54]    | 1.03    | [0.7-1.51]     | 1.03    | [0.75-1.41] |
| 6822 | Policeman   | 14/18              | 0.64     | [0.3-1.38]     | 0.81    | [0.56-1.18]    | 0.82    | [0.57-1.17]    | 0.83    | [0.59-1.17]    | 0.86    | [0.64-1.15] |
| 6824 | Private police guard  | 5/9                | 0.91     | [0.27-3.09]    | 0.97    | [0.55-1.71]    | 0.98    | [0.57-1.68]    | 0.99    | [0.61-1.62]    | 1.01    | [0.7-1.46]  |
| 6894 | Watchman  | 11/8               | 1.72     | [0.63-4.66]    | 1.28    | [0.8-2.06]     | 1.27    | [0.8-2.01]     | 1.23    | [0.81-1.89]    | 1.16    | [0.82-1.62] |
| 6111 | General farmer  | 27/21              | 1.29     | [0.69-2.41]    | 1.13    | [0.84-1.54]    | 1.13    | [0.84-1.53]    | 1.13    | [0.84-1.51]    | 1.11    | [0.85-1.43] |
| 6122 | Field Crop Farmer   | 7/15               | 0.45     | [0.17-1.18]    | 0.7     | [0.44-1.11]    | 0.72    | [0.46-1.12]    | 0.76    | [0.5-1.15]     | 0.85    | [0.61-1.18] |
| 6210 | Farm worker (general)   | 24/24              | 1.03     | [0.56-1.89]    | 1.02    | [0.76-1.37]    | 1.02    | [0.76-1.37]    | 1.02    | [0.77-1.36]    | 1.02    | [0.8-1.32]  |
| 6211 | Farm helper (general)   | 31/33              | 0.78     | [0.44-1.39]    | 0.89    | [0.67-1.18]    | 0.9     | [0.68-1.18]    | 0.9     | [0.69-1.18]    | 0.93    | [0.73-1.18] |
| 6221 | Field Crop Farm Worker (General).   | 8/7                | 1.14     | [0.39-3.37]    | 1.07    | [0.64-1.78]    | 1.07    | [0.65-1.74]    | 1.06    | [0.68-1.67]    | 1.05    | [0.74-1.49] |
| 6229 | Other field crop and vegetable farm workers   | 8/8                | 1.21     | [0.38-3.87]    | 1.1     | [0.64-1.89]    | 1.09    | [0.65-1.84]    | 1.09    | [0.68-1.74]    | 1.06    | [0.74-1.52] |
| 6233 | Vineyard worker   | 6/5                | 1.27     | [0.37-4.28]    | 1.12    | [0.64-1.96]    | 1.11    | [0.65-1.91]    | 1.1     | [0.68-1.79]    | 1.07    | [0.74-1.54] |
| 6239 | Other orchard, vineyard and related tree and shrub crop workers   | 5/8                | 0.79     | [0.21-3.06]    | 0.92    | [0.5-1.7]      | 0.93    | [0.52-1.67]    | 0.96    | [0.57-1.61]    | 0.99    | [0.68-1.45] |
| 6242 | Beef cattle farm worker   | 5/5                | 1.01     | [0.29-3.59]    | 1.02    | [0.57-1.82]    | 1.02    | [0.59-1.78]    | 1.03    | [0.62-1.69]    | 1.03    | [0.71-1.49] |
| 6243 | Sheep farm worker   | 10/6               | 0.91     | [0.28-2.99]    | 0.97    | [0.56-1.69]    | 0.98    | [0.58-1.66]    | 0.99    | [0.61-1.6]     | 1.01    | [0.7-1.45]  |
| 6274 | Gardener  | 6/3                | 2.77     | [0.64-11.97]   | 1.53    | [0.79-2.95]    | 1.47    | [0.79-2.72]    | 1.36    | [0.79-2.33]    | 1.19    | [0.8-1.75]  |
| 6001 | Production supervisor and general foreman   | 5/7                | 0.68     | [0.17-2.69]    | 0.85    | [0.45-1.58]    | 0.86    | [0.47-1.54]    | 0.87    | [0.52-1.47]    | 0.9     | [0.61-1.32] |
| 6003 | Supervisor and general foreman.   | 5/2                | 2.38     | [0.39-14.6]    | 1.42    | [0.66-3.08]    | 1.38    | [0.68-2.8]     | 1.31    | [0.72-2.38]    | 1.22    | [0.81-1.83] |
| 6005 | Supervisor and general foreman. Manufac. of machinery and metal products  | 17/22              | 0.81     | [0.41-1.59]    | 0.91    | [0.66-1.26]    | 0.91    | [0.66-1.26]    | 0.93    | [0.68-1.26]    | 0.96    | [0.73-1.26] |
| 7006 | Supervisor and general foreman. fabrication products of paper, plastics, rubber, chemical and synthetic materials | 5/2                | 3.05     | [0.56-16.69]   | 1.55    | [0.74-3.25]    | 1.47    | [0.74-2.92]    | 1.34    | [0.75-2.41]    | 1.17    | [0.78-1.75] |
| 7007 | Supervisor and general foreman. Production of textiles and clothing manufacturing.                                | 6/10               | 0.55     | [0.19-1.6]     | 0.78    | [0.47-1.28]    | 0.79    | [0.49-1.29]    | 0.83    | [0.53-1.29]    | 0.91    | [0.64-1.28] |
| 7009 | Other production supervisors and general foremen  | 9/6                | 1.13     | [0.38-3.35]    | 1.05    | [0.63-1.74]    | 1.04    | [0.63-1.69]    | 1.02    | [0.65-1.6]     | 0.98    | [0.69-1.39] |
| 7111 | Quarryman   | 6/4                | 1.41     | [0.36-5.51]    | 1.15    | [0.62-2.13]    | 1.14    | [0.64-2.03]    | 1.11    | [0.67-1.85]    | 1.07    | [0.73-1.55] |

| ISCO | Label  | Cases/<br>controls | ML<br>OR [95% CI] | HR $\tau=0.76$   |                  | HR $\tau=0.59$   |                  | HR $\tau=0.41$   |                  | HR $\tau=0.23$   |                  |
|------|--|--------------------|-------------------|------------------|------------------|------------------|------------------|------------------|------------------|------------------|------------------|
|      |  |                    |                   | OR               | [95% CI]         | OR               | [95% CI]         | OR               | [95% CI]         | OR               | [95% CI]         |
| 2219 | Other metal smelting, converting and refining furnacemen               | 8/9                | 0.95 [0.33-2.75]  | 0.97 [0.57-1.66] | 0.97 [0.57-1.66] | 0.97 [0.57-1.66] | 0.97 [0.57-1.66] | 0.97 [0.57-1.66] | 0.97 [0.57-1.66] | 0.97 [0.57-1.66] | 0.97 [0.57-1.66] |
| 7222 | Hot-roller (steel)   | 6/5                | 1.48 [0.43-5.13]  | 1.2 [0.67-2.12]  | 1.18 [0.68-2.04] | 1.15 [0.7-1.89]  | 1.1 [0.75-1.6]   | 1.1 [0.75-1.6]   | 1.1 [0.75-1.6]   | 1.1 [0.75-1.6]   | 1.1 [0.75-1.6]   |
| 7242 | Metal Pourer.  | 6/5                | 1.15 [0.33-4.1]   | 1.09 [0.61-1.95] | 1.1 [0.63-1.91]  | 1.11 [0.68-1.83] | 1.13 [0.78-1.63] | 1.13 [0.78-1.63] | 1.13 [0.78-1.63] | 1.13 [0.78-1.63] | 1.13 [0.78-1.63] |
| 7414 | Mixing- and blending-machine operator (chemical and related processes) | 5/6                | 1.11 [0.32-3.88]  | 1.08 [0.6-1.93]  | 1.08 [0.62-1.91] | 1.1 [0.65-1.86]  | 1.11 [0.71-1.74] | 1.11 [0.71-1.74] | 1.11 [0.71-1.74] | 1.11 [0.71-1.74] | 1.11 [0.71-1.74] |
| 7733 | Meat cutter  | 7/4                | 2.25 [0.59-8.54]  | 1.42 [0.77-2.61] | 1.38 [0.77-2.46] | 1.3 [0.78-2.18]  | 1.17 [0.8-1.7]   | 1.17 [0.8-1.7]   | 1.17 [0.8-1.7]   | 1.17 [0.8-1.7]   | 1.17 [0.8-1.7]   |
| 7761 | Baker, General   | 8/3                | 2.33 [0.58-9.3]   | 1.43 [0.77-2.69] | 1.39 [0.77-2.52] | 1.31 [0.77-2.21] | 1.17 [0.8-1.71]  | 1.17 [0.8-1.71]  | 1.17 [0.8-1.71]  | 1.17 [0.8-1.71]  | 1.17 [0.8-1.71]  |
| 7912 | Tailor (made-to-measure garments)                                      | 13/10              | 1.45 [0.6-3.52]   | 1.19 [0.78-1.82] | 1.19 [0.78-1.79] | 1.17 [0.79-1.72] | 1.12 [0.82-1.54] | 1.12 [0.82-1.54] | 1.12 [0.82-1.54] | 1.12 [0.82-1.54] | 1.12 [0.82-1.54] |
| 7963 | Vehicle upholsterer  | 7/6                | 0.78 [0.24-2.54]  | 0.91 [0.52-1.57] | 0.92 [0.54-1.56] | 0.94 [0.58-1.52] | 0.98 [0.68-1.41] | 0.98 [0.68-1.41] | 0.98 [0.68-1.41] | 0.98 [0.68-1.41] | 0.98 [0.68-1.41] |
| 8112 | Cabinetmaker   | 9/14               | 0.71 [0.29-1.73]  | 0.86 [0.56-1.32] | 0.87 [0.57-1.32] | 0.89 [0.6-1.31]  | 0.94 [0.68-1.29] | 0.94 [0.68-1.29] | 0.94 [0.68-1.29] | 0.94 [0.68-1.29] | 0.94 [0.68-1.29] |
| 8311 | Blacksmith   | 7/6                | 1.33 [0.46-3.81]  | 1.16 [0.71-1.9]  | 1.16 [0.72-1.87] | 1.16 [0.75-1.79] | 1.16 [0.82-1.63] | 1.16 [0.82-1.63] | 1.16 [0.82-1.63] | 1.16 [0.82-1.63] | 1.16 [0.82-1.63] |
| 8322 | Tool and die maker   | 7/1                | 6.03 [1.19-30.63] | 2.04 [1-4.15]    | 1.89 [0.97-3.65] | 1.65 [0.93-2.91] | 1.36 [0.91-2.02] | 1.36 [0.91-2.02] | 1.36 [0.91-2.02] | 1.36 [0.91-2.02] | 1.36 [0.91-2.02] |
| 8332 | Lathe setter-operator  | 6/5                | 0.94 [0.26-3.32]  | 0.98 [0.55-1.76] | 0.99 [0.57-1.73] | 1 [0.61-1.65]    | 1.02 [0.7-1.47]  | 1.02 [0.7-1.47]  | 1.02 [0.7-1.47]  | 1.02 [0.7-1.47]  | 1.02 [0.7-1.47]  |
| 8341 | Machine-tool operator  | 32/31              | 1.29 [0.75-2.22]  | 1.14 [0.87-1.49] | 1.14 [0.88-1.48] | 1.14 [0.88-1.47] | 1.14 [0.91-1.44] | 1.14 [0.91-1.44] | 1.14 [0.91-1.44] | 1.14 [0.91-1.44] | 1.14 [0.91-1.44] |
| 8342 | Lathe operator   | 50/34              | 1.77 [1.09-2.89]  | 1.33 [1.04-1.69] | 1.32 [1.04-1.68] | 1.31 [1.04-1.66] | 1.28 [1.04-1.59] | 1.28 [1.04-1.59] | 1.28 [1.04-1.59] | 1.28 [1.04-1.59] | 1.28 [1.04-1.59] |
| 8343 | Milling-machine operator   | 13/13              | 1.07 [0.47-2.45]  | 1.05 [0.7-1.56]  | 1.05 [0.71-1.55] | 1.06 [0.74-1.54] | 1.09 [0.8-1.48]  | 1.09 [0.8-1.48]  | 1.09 [0.8-1.48]  | 1.09 [0.8-1.48]  | 1.09 [0.8-1.48]  |
| 8346 | Drilling-machine operator  | 16/22              | 0.88 [0.44-1.76]  | 0.95 [0.68-1.33] | 0.96 [0.69-1.33] | 0.98 [0.71-1.34] | 1.02 [0.77-1.34] | 1.02 [0.77-1.34] | 1.02 [0.77-1.34] | 1.02 [0.77-1.34] | 1.02 [0.77-1.34] |
| 8352 | Buffing- and Polishing-Machine Operator                                | 13/13              | 0.97 [0.43-2.16]  | 1 [0.68-1.47]    | 1 [0.69-1.47]    | 1.02 [0.71-1.46] | 1.06 [0.78-1.43] | 1.06 [0.78-1.43] | 1.06 [0.78-1.43] | 1.06 [0.78-1.43] | 1.06 [0.78-1.43] |
| 8392 | Gunsmith   | 8/7                | 1.14 [0.38-3.47]  | 1.09 [0.65-1.82] | 1.1 [0.67-1.8]   | 1.11 [0.71-1.74] | 1.13 [0.8-1.6]   | 1.13 [0.8-1.6]   | 1.13 [0.8-1.6]   | 1.13 [0.8-1.6]   | 1.13 [0.8-1.6]   |
| 8395 | Metal former (hand)  | 6/6                | 1.51 [0.44-5.22]  | 1.22 [0.69-2.16] | 1.22 [0.71-2.1]  | 1.21 [0.74-1.96] | 1.18 [0.82-1.7]  | 1.18 [0.82-1.7]  | 1.18 [0.82-1.7]  | 1.18 [0.82-1.7]  | 1.18 [0.82-1.7]  |
| 8396 | Metal-press operator   | 22/16              | 1.27 [0.61-2.67]  | 1.13 [0.79-1.62] | 1.13 [0.8-1.61]  | 1.14 [0.81-1.59] | 1.14 [0.86-1.52] | 1.14 [0.86-1.52] | 1.14 [0.86-1.52] | 1.14 [0.86-1.52] | 1.14 [0.86-1.52] |
| 8398 | Power-Shear Operator.  | 5/8                | 0.51 [0.16-1.61]  | 0.75 [0.43-1.28] | 0.77 [0.45-1.29] | 0.81 [0.5-1.3]   | 0.89 [0.61-1.29] | 0.89 [0.61-1.29] | 0.89 [0.61-1.29] | 0.89 [0.61-1.29] | 0.89 [0.61-1.29] |
| 8399 | Other blacksmiths, toolmakers and machine-tool operator                | 29/25              | 1.31 [0.7-2.45]   | 1.15 [0.84-1.56] | 1.15 [0.85-1.55] | 1.15 [0.86-1.54] | 1.15 [0.89-1.49] | 1.15 [0.89-1.49] | 1.15 [0.89-1.49] | 1.15 [0.89-1.49] | 1.15 [0.89-1.49] |
| 8410 | Machinery fitter   | 5/1                | 7.63 [0.83-69.9]  | 1.93 [0.79-4.69] | 1.71 [0.77-3.81] | 1.44 [0.75-2.78] | 1.19 [0.77-1.83] | 1.19 [0.77-1.83] | 1.19 [0.77-1.83] | 1.19 [0.77-1.83] | 1.19 [0.77-1.83] |
| 8411 | Machinery fitter-assembler   | 14/23              | 0.89 [0.44-1.78]  | 0.95 [0.68-1.33] | 0.95 [0.68-1.33] | 0.96 [0.7-1.33]  | 0.98 [0.74-1.3]  | 0.98 [0.74-1.3]  | 0.98 [0.74-1.3]  | 0.98 [0.74-1.3]  | 0.98 [0.74-1.3]  |
| 8419 | Other machinery fitters and machine assemblers                         | 8/10               | 0.96 [0.35-2.67]  | 0.99 [0.61-1.61] | 1 [0.63-1.59]    | 1.01 [0.65-1.55] | 1.02 [0.72-1.44] | 1.02 [0.72-1.44] | 1.02 [0.72-1.44] | 1.02 [0.72-1.44] | 1.02 [0.72-1.44] |
| 8432 | Automobile mechanic  | 20/16              | 1.69 [0.8-3.56]   | 1.28 [0.89-1.83] | 1.26 [0.89-1.79] | 1.23 [0.89-1.72] | 1.16 [0.88-1.53] | 1.16 [0.88-1.53] | 1.16 [0.88-1.53] | 1.16 [0.88-1.53] | 1.16 [0.88-1.53] |
| 8433 | Motor-truck mechanic   | 6/6                | 0.94 [0.28-3.18]  | 0.99 [0.57-1.71] | 0.99 [0.59-1.67] | 1 [0.63-1.6]     | 1.02 [0.72-1.43] | 1.02 [0.72-1.43] | 1.02 [0.72-1.43] | 1.02 [0.72-1.43] | 1.02 [0.72-1.43] |
| 8439 | Other motor-vehicle mechanics  | 8/9                | 0.88 [0.27-2.86]  | 0.96 [0.56-1.64] | 0.97 [0.58-1.61] | 0.98 [0.62-1.55] | 1 [0.71-1.41]    | 1 [0.71-1.41]    | 1 [0.71-1.41]    | 1 [0.71-1.41]    | 1 [0.71-1.41]    |
| 8491 | Machinery mechanic (general)   | 17/16              | 1 [0.48-2.1]      | 1 [0.7-1.44]     | 1.01 [0.71-1.43] | 1.01 [0.73-1.4]  | 1.02 [0.77-1.34] | 1.02 [0.77-1.34] | 1.02 [0.77-1.34] | 1.02 [0.77-1.34] | 1.02 [0.77-1.34] |
| 8493 | Metalworking machine-tool mechanic                                     | 13/8               | 2.31 [0.88-6.06]  | 1.45 [0.92-2.28] | 1.41 [0.91-2.18] | 1.34 [0.89-1.99] | 1.19 [0.87-1.64] | 1.19 [0.87-1.64] | 1.19 [0.87-1.64] | 1.19 [0.87-1.64] | 1.19 [0.87-1.64] |

| ISCO | Label   | Cases/<br>controls | ML<br>OR [95% CI] | HR $\tau=0.76$   |                  | HR $\tau=0.59$   |                  | HR $\tau=0.41$ |          | HR $\tau=0.23$ |          |
|------|---|--------------------|-------------------|------------------|------------------|------------------|------------------|----------------|----------|----------------|----------|
|      |   |                    |                   | OR               | [95% CI]         | OR               | [95% CI]         | OR             | [95% CI] | OR             | [95% CI] |
| 8494 | Printing machinery mechanic   | 6/6                | 0.89 [0.3-2.67]   | 0.96 [0.58-1.59] | 0.97 [0.6-1.57]  | 0.98 [0.64-1.52] | 1                | [0.72-1.4]     |          |                |          |
| 8497 | Plant maintenance mechanic  | 8/18               | 0.61 [0.25-1.49]  | 0.8 [0.52-1.22]  | 0.8 [0.53-1.21]  | 0.82 [0.56-1.21] | 0.86 [0.62-1.18] |                |          |                |          |
| 8498 | Mechanical products inspector and tester                                    | 16/32              | 0.54 [0.29-1.01]  | 0.75 [0.56-1.02] | 0.76 [0.57-1.02] | 0.78 [0.59-1.04] | 0.84 [0.66-1.09] |                |          |                |          |
| 8499 | Other machinery fitters, machine assemblers and precision-instrument makers | 46/38              | 1.36 [0.84-2.21]  | 1.16 [0.92-1.48] | 1.16 [0.92-1.47] | 1.16 [0.92-1.46] | 1.14 [0.91-1.41] |                |          |                |          |
| 8532 | Electrical equipment assembler  | 5/5                | 0.47 [0.1-2.09]   | 0.75 [0.38-1.46] | 0.78 [0.41-1.47] | 0.84 [0.48-1.45] | 0.93 [0.63-1.38] |                |          |                |          |
| 8551 | Electrician, general  | 16/12              | 1.47 [0.6-3.57]   | 1.2 [0.78-1.84]  | 1.19 [0.79-1.8]  | 1.17 [0.79-1.73] | 1.12 [0.82-1.55] |                |          |                |          |
| 8552 | Building electrician  | 5/7                | 0.65 [0.17-2.43]  | 0.84 [0.46-1.54] | 0.86 [0.49-1.53] | 0.9 [0.54-1.5]   | 0.96 [0.66-1.4]  |                |          |                |          |
| 8556 | Maintenance electrician   | 10/11              | 1.41 [0.56-3.52]  | 1.18 [0.76-1.82] | 1.17 [0.76-1.79] | 1.15 [0.77-1.72] | 1.11 [0.8-1.54]  |                |          |                |          |
| 8710 | Plumber (general)   | 12/11              | 1 [0.4-2.52]      | 1 [0.64-1.55]    | 1 [0.65-1.54]    | 1 [0.67-1.5]     | 1 [0.71-1.4]     |                |          |                |          |
| 8721 | Gas and electric welder (general)   | 29/20              | 1.21 [0.64-2.31]  | 1.09 [0.8-1.5]   | 1.09 [0.8-1.49]  | 1.08 [0.8-1.46]  | 1.06 [0.81-1.39] |                |          |                |          |
| 8722 | Electric arc welder (hand)  | 6/4                | 1.69 [0.45-6.33]  | 1.24 [0.67-2.27] | 1.21 [0.68-2.15] | 1.16 [0.69-1.94] | 1.08 [0.73-1.6]  |                |          |                |          |
| 8737 | Vehicle sheet-metal worker  | 15/14              | 1.25 [0.55-2.86]  | 1.11 [0.74-1.65] | 1.1 [0.75-1.63]  | 1.09 [0.75-1.58] | 1.06 [0.77-1.46] |                |          |                |          |
| 8743 | Structural steel worker (workshop)  | 13/11              | 1.17 [0.5-2.76]   | 1.07 [0.71-1.62] | 1.07 [0.71-1.6]  | 1.06 [0.72-1.55] | 1.04 [0.75-1.44] |                |          |                |          |
| 8744 | Constructional steel erector  | 10/14              | 0.58 [0.21-1.6]   | 0.78 [0.48-1.27] | 0.8 [0.5-1.27]   | 0.83 [0.54-1.28] | 0.89 [0.63-1.27] |                |          |                |          |
| 9013 | Rubber extruding-machine operator   | 5/1                | 5.07 [0.43-59.92] | 1.59 [0.61-4.19] | 1.44 [0.6-3.43]  | 1.25 [0.61-2.56] | 1.07 [0.64-1.8]  |                |          |                |          |
| 9015 | Plastics injection-moulding-machine operator                                | 5/4                | 1.58 [0.4-6.32]   | 1.24 [0.66-2.36] | 1.24 [0.67-2.29] | 1.21 [0.69-2.13] | 1.17 [0.73-1.86] |                |          |                |          |
| 9019 | Other rubber and plastics products makers                                   | 5/0                | NA                | 0.97 [0.24-4.81] | 1.07 [0.33-3.43] | 1.06 [0.47-2.38] | 1.04 [0.66-1.65] |                |          |                |          |
| 9022 | Tire builder  | 6/3                | 0.85 [0.16-4.45]  | 0.97 [0.47-2]    | 0.97 [0.49-1.94] | 0.98 [0.53-1.8]  | 0.98 [0.61-1.58] |                |          |                |          |
| 9211 | Printer, general  | 6/9                | 0.38 [0.11-1.33]  | 0.66 [0.37-1.19] | 0.69 [0.39-1.2]  | 0.73 [0.44-1.22] | 0.8 [0.53-1.21]  |                |          |                |          |
| 9312 | Building painter  | 21/18              | 1 [0.51-1.97]     | 1 [0.72-1.39]    | 1 [0.72-1.38]    | 0.99 [0.72-1.36] | 0.96 [0.72-1.29] |                |          |                |          |
| 9392 | Brush-Painter (except Construction)   | 5/3                | 4 [0.7-22.81]     | 1.63 [0.76-3.5]  | 1.5 [0.73-3.05]  | 1.29 [0.7-2.4]   | 1.06 [0.67-1.67] |                |          |                |          |
| 9393 | Spray-Painter (except Construction)   | 9/0                | NA                | NA               | NA               | NA               | NA               |                |          |                |          |
| 9396 | Automobile painter  | 9/14               | 0.7 [0.26-1.88]   | 0.86 [0.54-1.38] | 0.88 [0.55-1.39] | 0.89 [0.58-1.38] | 0.93 [0.63-1.37] |                |          |                |          |
| 9399 | Other painters  | 5/11               | 0.55 [0.2-1.54]   | 0.76 [0.47-1.25] | 0.78 [0.48-1.25] | 0.8 [0.51-1.25]  | 0.84 [0.57-1.22] |                |          |                |          |
| 9498 | Quality inspector   | 7/13               | 0.59 [0.22-1.6]   | 0.8 [0.5-1.28]   | 0.81 [0.51-1.28] | 0.84 [0.55-1.28] | 0.91 [0.65-1.27] |                |          |                |          |
| 9511 | Bricklayers, Stonemasons and Tile Setters                                   | 27/20              | 1.23 [0.66-2.32]  | 1.11 [0.81-1.51] | 1.11 [0.82-1.5]  | 1.1 [0.82-1.48]  | 1.09 [0.84-1.41] |                |          |                |          |
| 9512 | Bricklayer (construction)   | 64/90              | 0.72 [0.5-1.04]   | 0.85 [0.71-1.02] | 0.85 [0.71-1.02] | 0.86 [0.72-1.03] | 0.87 [0.74-1.04] |                |          |                |          |
| 9541 | Carpenter, General  | 5/16               | 0.22 [0.07-0.68]  | 0.52 [0.31-0.89] | 0.55 [0.33-0.92] | 0.61 [0.39-0.98] | 0.77 [0.54-1.09] |                |          |                |          |
| 9547 | Bench carpenter   | 14/11              | 1.28 [0.54-3.03]  | 1.13 [0.75-1.7]  | 1.13 [0.75-1.68] | 1.12 [0.76-1.63] | 1.09 [0.8-1.49]  |                |          |                |          |
| 9551 | Plasterer, general  | 5/2                | 2.25 [0.41-12.2]  | 1.38 [0.66-2.89] | 1.33 [0.67-2.64] | 1.25 [0.7-2.24]  | 1.13 [0.76-1.69] |                |          |                |          |
| 9713 | Railway and road vehicle loader   | 5/3                | 1 [0.19-5.34]     | 0.99 [0.48-2.04] | 0.98 [0.5-1.92]  | 0.97 [0.54-1.72] | 0.94 [0.63-1.41] |                |          |                |          |
| 9714 | Boat loader (liquids and gases)   | 37/43              | 0.97 [0.6-1.57]   | 0.99 [0.78-1.25] | 0.99 [0.78-1.25] | 0.99 [0.79-1.25] | 1 [0.81-1.24]    |                |          |                |          |
| 9732 | Boat loader (liquids and gases)   | 10/7               | 1.74 [0.59-5.12]  | 1.29 [0.77-2.14] | 1.27 [0.78-2.07] | 1.23 [0.79-1.93] | 1.15 [0.81-1.63] |                |          |                |          |
| 9792 | Lifting-truck operator  | 12/19              | 0.78 [0.38-1.61]  | 0.89 [0.62-1.26] | 0.89 [0.63-1.26] | 0.89 [0.64-1.24] | 0.9 [0.68-1.2]   |                |          |                |          |

| ISCO | Label   | Cases/<br>controls | ML<br>OR [95% CI] | HR $\tau=0.76$   |                  | HR $\tau=0.59$   |                  | HR $\tau=0.41$ |          | HR $\tau=0.23$ |          |
|------|---|--------------------|-------------------|------------------|------------------|------------------|------------------|----------------|----------|----------------|----------|
|      |   |                    |                   | OR               | [95% CI]         | OR               | [95% CI]         | OR             | [95% CI] | OR             | [95% CI] |
| 9854 | Motor bus driver                                  | 6/14               | 0.64 [0.23-1.81]  | 0.82 [0.5-1.33]  | 0.82 [0.51-1.32] | 0.84 [0.54-1.3]  | 0.87 [0.62-1.24] |                |          |                |          |
| 9855 | Lorry and van driver (local transport)            | 27/31              | 0.81 [0.47-1.4]   | 0.9 [0.69-1.18]  | 0.9 [0.69-1.18]  | 0.91 [0.7-1.18]  | 0.91 [0.71-1.18] |                |          |                |          |
| 9856 | Lorry and van driver<br>(long-distance transport) | 25/17              | 1.06 [0.54-2.08]  | 1.03 [0.74-1.42] | 1.02 [0.74-1.41] | 1.02 [0.74-1.38] | 0.99 [0.75-1.3]  |                |          |                |          |
| 9991 | Labourer  | 82/85              | 0.86 [0.6-1.23]   | 0.93 [0.78-1.11] | 0.93 [0.78-1.11] | 0.93 [0.78-1.11] | 0.94 [0.8-1.11]  |                |          |                |          |

**Supplementary Table 4. Results for the risk of BC due to occupations performed longest in life with at least 5 cases. Models are adjusted for age, smoking status, intensity of smoking and study.**

| IACO Label  | Cases/<br>controls | ML<br>OR [95% CI] | HR $\tau=0.76$ |             | HR $\tau=0.59$ |             | HR $\tau=0.41$ |             | HR $\tau=0.23$ |             |
|---|--------------------|-------------------|----------------|-------------|----------------|-------------|----------------|-------------|----------------|-------------|
|   |                    |                   | OR             | [95% CI]    | OR             | [95% CI]    | OR             | [95% CI]    | OR             | [95% CI]    |
| 0022 Mechanical draughtsman   | 9/7                | 1.99 [0.65-6.1]   | 1.39           | [0.82-2.34] | 1.37           | [0.83-2.27] | 1.33           | [0.84-2.12] | 1.25           | [0.87-1.79] |
| 0099 Other Engineering Technicians.   | 5/3                | 1.48 [0.3-7.17]   | 1.21           | [0.6-2.45]  | 1.21           | [0.63-2.33] | 1.2            | [0.68-2.12] | 1.16           | [0.78-1.74] |
| 2011 General Manager.   | 16/17              | 1.17 [0.55-2.48]  | 1.09           | [0.75-1.57] | 1.09           | [0.76-1.56] | 1.1            | [0.78-1.55] | 1.11           | [0.83-1.49] |
| 2021 Production Manager (except Farm).  | 7/1                | 6.41 [0.75-54.97] | 1.97           | [0.82-4.73] | 1.79           | [0.81-3.97] | 1.55           | [0.81-2.98] | 1.29           | [0.84-1.99] |
| 2199 Other Managers.  | 5/8                | 0.44 [0.08-2.3]   | 0.76           | [0.37-1.58] | 0.81           | [0.41-1.6]  | 0.89           | [0.5-1.6]   | 1.01           | [0.67-1.52] |
| 3001 Government Executive Official.   | 5/5                | 0.66 [0.18-2.36]  | 0.86           | [0.48-1.55] | 0.89           | [0.51-1.56] | 0.94           | [0.57-1.55] | 1.02           | [0.7-1.49]  |
| 3703 postman  | 5/0                | NA                | 1.21           | [0.27-5.53] | 1.2            | [0.37-3.91] | 1.18           | [0.52-2.69] | 1.15           | [0.72-1.84] |
| 3922 Material planning clerk  | 6/5                | 1.85 [0.53-6.4]   | 1.34           | [0.75-2.38] | 1.32           | [0.76-2.29] | 1.29           | [0.78-2.12] | 1.22           | [0.84-1.77] |
| 3931 Office clerk   | 34/23              | 1.57 [0.86-2.89]  | 1.25           | [0.93-1.69] | 1.25           | [0.93-1.68] | 1.25           | [0.94-1.65] | 1.22           | [0.95-1.57] |
| 4002 Working Proprietor (Wholesale Trade).                                    | 6/5                | 1.29 [0.38-4.4]   | 1.15           | [0.65-2.02] | 1.15           | [0.67-1.98] | 1.15           | [0.7-1.88]  | 1.14           | [0.79-1.66] |
| 4003 Working proprietor (retail trade)  | 14/11              | 1.2 [0.51-2.83]   | 1.1            | [0.73-1.67] | 1.11           | [0.74-1.66] | 1.11           | [0.76-1.63] | 1.12           | [0.82-1.54] |
| 4019 Other Sales Supervisors  | 5/2                | 2.51 [0.41-15.32] | 1.47           | [0.68-3.21] | 1.43           | [0.7-2.92]  | 1.34           | [0.73-2.47] | 1.22           | [0.81-1.86] |
| 4022 Commercial traveller   | 9/16               | 0.48 [0.2-1.16]   | 0.73           | [0.48-1.11] | 0.74           | [0.49-1.12] | 0.78           | [0.53-1.16] | 0.89           | [0.64-1.22] |
| 4023 Manufacturers' agent   | 9/9                | 0.92 [0.3-2.8]    | 0.99           | [0.58-1.66] | 1              | [0.6-1.66]  | 1.02           | [0.65-1.63] | 1.07           | [0.75-1.53] |
| 4013 Cook, except private service   | 9/5                | 2.57 [0.81-8.14]  | 1.55           | [0.9-2.65]  | 1.51           | [0.9-2.54]  | 1.45           | [0.9-2.32]  | 1.3            | [0.91-1.88] |
| 4022 Street vendor  | 5/9                | 0.47 [0.15-1.47]  | 0.74           | [0.43-1.26] | 0.76           | [0.46-1.28] | 0.82           | [0.51-1.31] | 0.94           | [0.65-1.35] |
| 5025 Bartender  | 5/1                | 5.6 [0.63-50.03]  | 1.87           | [0.77-4.55] | 1.72           | [0.77-3.83] | 1.51           | [0.78-2.91] | 1.28           | [0.83-1.97] |
| 5022 Policeman  | 8/10               | 0.75 [0.27-2.05]  | 0.9            | [0.56-1.46] | 0.92           | [0.58-1.47] | 0.96           | [0.62-1.49] | 1.03           | [0.71-1.5]  |
| 6011 General farmer   | 12/13              | 0.92 [0.39-2.15]  | 0.98           | [0.65-1.47] | 0.98           | [0.66-1.47] | 1              | [0.69-1.46] | 1.04           | [0.76-1.43] |
| 6010 Farm worker  | 10/6               | 2 [0.67-5.99]     | 1.39           | [0.83-2.33] | 1.37           | [0.84-2.26] | 1.34           | [0.85-2.11] | 1.25           | [0.87-1.79] |
| Supervisor and general foreman, Manufacturing of machinery and metal products | 11/9               | 1.32 [0.51-3.42]  | 1.16           | [0.73-1.83] | 1.16           | [0.74-1.81] | 1.16           | [0.76-1.77] | 1.14           | [0.79-1.66] |
| Other production supervisors and general foremen                              | 5/2                | 1.77 [0.31-10.1]  | 1.3            | [0.61-2.79] | 1.29           | [0.63-2.62] | 1.25           | [0.67-2.33] | 1.19           | [0.75-1.89] |
| 7033 Meat Cutter  | 5/3                | 2.04 [0.41-10.05] | 1.38           | [0.68-2.79] | 1.35           | [0.7-2.61]  | 1.3            | [0.73-2.3]  | 1.21           | [0.81-1.82] |
| 7061 Baker, General   | 5/2                | 2.62 [0.48-14.29] | 1.51           | [0.72-3.16] | 1.46           | [0.73-2.9]  | 1.37           | [0.76-2.48] | 1.24           | [0.82-1.87] |
| 7012 Tailor (made-to-measure garments)  | 6/2                | 2.62 [0.5-13.85]  | 1.51           | [0.73-3.14] | 1.46           | [0.74-2.89] | 1.37           | [0.76-2.47] | 1.24           | [0.82-1.87] |
| 8341 Machine-tool operator  | 9/12               | 1.01 [0.4-2.54]   | 1.02           | [0.65-1.59] | 1.03           | [0.67-1.58] | 1.05           | [0.7-1.57]  | 1.09           | [0.77-1.52] |

| ISCO Label  | Cases/<br>controls | ML<br>OR [95% CI] | HR $\tau=0.76$ |             | HR $\tau=0.59$ |             | HR $\tau=0.41$ |             | HR $\tau=0.23$ |             |
|---|--------------------|-------------------|----------------|-------------|----------------|-------------|----------------|-------------|----------------|-------------|
|   |                    |                   | OR             | [95% CI]    | OR             | [95% CI]    | OR             | [95% CI]    | OR             | [95% CI]    |
| 8442 Lathe operator   | 20/13              | 2.36 [1.06-5.28]  | 1.5            | [1.02-2.22] | 1.49           | [1.02-2.17] | 1.44           | [1.01-2.07] | 1.34           | [0.98-1.83] |
| 8443 Milling-machine operator   | 8/6                | 1.54 [0.49-4.84]  | 1.23           | [0.72-2.1]  | 1.23           | [0.73-2.05] | 1.21           | [0.76-1.94] | 1.19           | [0.82-1.72] |
| 8446 Drilling-machine operator  | 6/7                | 1.28 [0.4-4.11]   | 1.14           | [0.66-1.96] | 1.14           | [0.68-1.92] | 1.14           | [0.71-1.84] | 1.15           | [0.79-1.67] |
| 8452 Buffing- and Polishing-Machine Operator  | 6/5                | 1.8 [0.52-6.29]   | 1.31           | [0.74-2.34] | 1.3            | [0.75-2.25] | 1.27           | [0.77-2.08] | 1.21           | [0.82-1.78] |
| 8496 Metal-press operator   | 7/10               | 0.8 [0.26-2.47]   | 0.93           | [0.55-1.57] | 0.95           | [0.57-1.57] | 0.98           | [0.62-1.57] | 1.06           | [0.73-1.53] |
| 8499 Other blacksmiths. toolmakers<br>and machine-tool operator                     | 11/10              | 1.02 [0.36-2.91]  | 1.03           | [0.63-1.69] | 1.04           | [0.65-1.68] | 1.06           | [0.68-1.65] | 1.1            | [0.77-1.57] |
| 8492 Automobile mechanic  | 7/3                | 2.4 [0.52-11.1]   | 1.44           | [0.74-2.81] | 1.39           | [0.74-2.61] | 1.32           | [0.76-2.28] | 1.22           | [0.8-1.86]  |
| 8491 Machinery mechanic (general)   | 7/9                | 0.89 [0.3-2.68]   | 0.98           | [0.59-1.64] | 1              | [0.61-1.64] | 1.03           | [0.66-1.63] | 1.09           | [0.75-1.59] |
| 8493 Metalworking machine-tool mechanic   | 6/3                | 3.52 [0.76-16.37] | 1.65           | [0.84-3.22] | 1.56           | [0.83-2.92] | 1.43           | [0.82-2.47] | 1.27           | [0.83-1.93] |
| 8498 Mechanical products inspector and tester                                       | 7/10               | 0.84 [0.3-2.34]   | 0.96           | [0.59-1.54] | 0.97           | [0.61-1.55] | 1.01           | [0.66-1.55] | 1.07           | [0.74-1.54] |
| 8499 Other machinery fitters. machine assemblers and<br>precision-instrument makers | 18/13              | 1.28 [0.58-2.79]  | 1.14           | [0.78-1.66] | 1.14           | [0.78-1.66] | 1.14           | [0.79-1.64] | 1.13           | [0.81-1.58] |
| 8451 Electrician. general   | 7/5                | 3 [0.68-13.27]    | 1.61           | [0.83-3.15] | 1.56           | [0.83-2.92] | 1.45           | [0.84-2.52] | 1.28           | [0.86-1.91] |
| 8410 Plumber  | 6/7                | 0.75 [0.23-2.43]  | 0.87           | [0.49-1.53] | 0.87           | [0.5-1.51]  | 0.87           | [0.51-1.48] | 0.88           | [0.55-1.41] |
| 8721 Gas and electric welder  | 8/8                | 0.81 [0.28-2.35]  | 0.9            | [0.54-1.5]  | 0.9            | [0.54-1.49] | 0.9            | [0.55-1.46] | 0.89           | [0.57-1.39] |
| 9022 Tire builder   | 6/1                | 2.58 [0.26-25.68] | 1.61           | [0.51-5.07] | 1.61           | [0.51-5.07] | 1.61           | [0.51-5.07] | 1.61           | [0.51-5.07] |
| 9112 Building painter   | 14/9               | 1.21 [0.48-3.03]  | 1.1            | [0.7-1.74]  | 1.1            | [0.7-1.74]  | 1.1            | [0.7-1.74]  | 1.1            | [0.7-1.74]  |
| 9112 Bricklayers. Stonemasons and Tile Setters                                      | 34/39              | 0.85 [0.5-1.43]   | 0.93           | [0.72-1.2]  | 0.93           | [0.72-1.21] | 0.94           | [0.73-1.21] | 0.97           | [0.77-1.22] |
| 9451 Plasterer. general   | 5/1                | 4.11 [0.46-36.81] | 1.69           | [0.7-4.12]  | 1.58           | [0.71-3.53] | 1.43           | [0.74-2.76] | 1.25           | [0.81-1.92] |
| 9414 Boat loader (liquids and gases)  | 16/19              | 1 [0.49-2.06]     | 1.01           | [0.71-1.43] | 1.02           | [0.72-1.43] | 1.03           | [0.74-1.43] | 1.06           | [0.79-1.41] |
| 9432 Bridge- or gantry-crane operator   | 6/2                | 4.35 [0.5-37.82]  | 1.73           | [0.72-4.17] | 1.61           | [0.73-3.58] | 1.44           | [0.75-2.78] | 1.25           | [0.82-1.93] |
| 9492 Lifting-truck operator   | 5/3                | 1.77 [0.33-9.41]  | 1.3            | [0.62-2.73] | 1.29           | [0.64-2.58] | 1.26           | [0.68-2.31] | 1.19           | [0.76-1.89] |
| 9855 Lorry and van driver (local transport)   | 12/12              | 0.93 [0.4-2.17]   | 0.97           | [0.63-1.47] | 0.97           | [0.63-1.47] | 0.97           | [0.63-1.47] | 0.97           | [0.63-1.47] |
| 9856 Lorry and van driver (long-distance transport)                                 | 16/7               | 1.85 [0.71-4.82]  | 1.35           | [0.85-2.13] | 1.34           | [0.86-2.09] | 1.31           | [0.86-2]    | 1.24           | [0.86-1.8]  |
| 9991 Labourer   | 10/9               | 1.22 [0.46-3.2]   | 1.11           | [0.7-1.77]  | 1.12           | [0.71-1.75] | 1.12           | [0.74-1.70] | 1.13           | [0.8-1.58]  |

**Supplementary Table 5. Results of the multivariable logistic regression models for risk of BC due to exposure (never/ever) and cumulative exposure (no/low/high) to carcinogens in high grade BC. Models are adjusted for age, smoking status, intensity of smoking and study**

| Carcinogen   | Exposure | Cases / Controls | OR [95% CI]      | Exposure | Cases / Controls | OR [95% CI]      |
|--|----------|------------------|------------------|----------|------------------|------------------|
| <b>Aliphatic and alicyclic hydrocarbon solvents (ALHC)</b> | Ever     | 40/119           | 0.78 [0.51-1.21] | High     | 25/60            | 1.00 [0.57-1.73] |
|  | Never    | 305/859          | Ref.             | Low      | 15/59            | 0.58 [0.3-1.11]  |
|  |          |                  |                  | No       | 305/859          | Ref.             |
| <b>Aromatic amines (AA)</b>                                | Ever     | 22/61            | 0 [0-0]          | High     | NA               | NA               |
|  | Never    | 320/913          | Ref.             | Low      | NA               | NA               |
|  |          |                  |                  | No       | NA               | Ref.             |
| <b>Aromatic hydrocarbon solvents (ARHC)</b>                | Ever     | 87/220           | 0.96 [0.69-1.32] | High     | 57/116           | 1.16 [0.78-1.73] |
|  | Never    | 258/758          | Ref.             | Low      | 30/104           | 0.73 [0.45-1.18] |
|  |          |                  |                  | No       | 258/758          | Ref.             |
| <b>Arsenic (AS)</b>  | Ever     | 30/81            | 0.85 [0.52-1.38] | High     | 17/41            | 0.83 [0.43-1.62] |
|  | Never    | 315/897          | Ref.             | Low      | 13/40            | 0.86 [0.44-1.71] |
|  |          |                  |                  | No       | 315/897          | Ref.             |
| <b>Benzene (BENZ)</b>                                      | Ever     | 68/187           | 0.94 [0.66-1.33] | High     | 33/94            | 0.8 [0.5-1.3]    |
|  | Never    | 277/791          | Ref.             | Low      | 35/93            | 1.09 [0.68-1.74] |
|  |          |                  |                  | No       | 277/791          | Ref.             |
| <b>Benzo(a)pyrene (BAP)</b>                                | Ever     | 111/294          | 1.06 [0.79-1.42] | High     | 55/149           | 0.92 [0.62-1.35] |
|  | Never    | 234/684          | Ref.             | Low      | 56/145           | 1.21 [0.83-1.77] |
|  |          |                  |                  | No       | 234/684          | Ref.             |
| <b>Cadmium (CD)</b>  | Ever     | 128/334          | 1.11 [0.84-1.48] | High     | 61/172           | 0.99 [0.68-1.43] |
|  | Never    | 217/644          | Ref.             | Low      | 67/162           | 1.25 [0.87-1.8]  |
|  |          |                  |                  | No       | 217/644          | Ref.             |
| <b>Chromium (CR)</b>                                       | Ever     | 166/430          | 1.14 [0.87-1.5]  | High     | 91/216           | 1.2 [0.86-1.68]  |
|  | Never    | 179/548          | Ref.             | Low      | 75/214           | 1.08 [0.76-1.53] |
|  |          |                  |                  | No       | 179/548          | Ref.             |
| <b>Detergents (DET)</b>                                    | Ever     | 185/508          | 1.06 [0.8-1.39]  | High     | 102/258          | 1.18 [0.85-1.63] |
|  | Never    | 160/470          | Ref.             | Low      | 83/250           | 0.93 [0.66-1.32] |
|  |          |                  |                  | No       | 160/470          | Ref.             |
| <b>Diesel engine exhaust (DEEX)</b>                        | Ever     | 82/256           | 0.81 [0.59-1.11] | High     | 52/129           | 0.99 [0.67-1.47] |
|  | Never    | 263/722          | Ref.             | Low      | 30/127           | 0.61 [0.38-0.98] |
|  |          |                  |                  | No       | 263/722          | Ref.             |
| <b>Fungicides (FUNG)</b>                                   | Ever     | 33/89            | 0.93 [0.58-1.49] | High     | 17/45            | 0.86 [0.45-1.64] |
|  | Never    | 312/889          | Ref.             | Low      | 16/44            | 1.01 [0.53-1.92] |
|  |          |                  |                  | No       | 312/889          | Ref.             |
| <b>Gasoline (GASO)</b>                                     | Ever     | 41/116           | 0.96 [0.63-1.47] | High     | 24/61            | 1 [0.57-1.75]    |
|  | Never    | 304/862          | Ref.             | Low      | 17/55            | 0.91 [0.49-1.69] |
|  |          |                  |                  | No       | 304/862          | Ref.             |
| <b>Gasoline engine exhaust (GEEX)</b>                      | Ever     | 93/267           | 0.84 [0.61-1.15] | High     | 55/146           | 0.91 [0.62-1.32] |
|  | Never    | 252/711          | Ref.             | Low      | 38/121           | 0.75 [0.47-1.18] |
|  |          |                  |                  | No       | 252/711          | Ref.             |
| <b>Herbicides (HERB)</b>                                   | Ever     | 32/83            | 0.95 [0.59-1.53] | High     | 16/45            | 0.75 [0.38-1.47] |
|  | Never    | 313/895          | Ref.             | Low      | 16/38            | 1.2 [0.63-2.3]   |
|  |          |                  |                  | No       | 313/895          | Ref.             |
| <b>Insecticides (INSC)</b>                                 | Ever     | 32/86            | 0.93 [0.58-1.5]  | High     | 15/44            | 0.79 [0.4-1.55]  |
|  | Never    | 313/892          | Ref.             | Low      | 17/42            | 1.09 [0.58-2.06] |
|  |          |                  |                  | No       | 313/892          | Ref.             |



| Cancerogen                                     | Exposure | Cases / Controls | OR [95% CI]      | Exposure | Cases / Controls | OR [95% CI]      |
|--|----------|------------------|------------------|----------|------------------|------------------|
| <b>Ionizing radiation (IRAD)</b>               | Ever     | 6/33             | 0.52 [0.2-1.4]   | High     | 6/18             | 0.94 [0.34-2.66] |
|  | Never    | 339/945          | Ref.             | Low      | 0/15             | 0 [0-0]          |
|  |          |                  |                  | No       | 339/945          | Ref.             |
| <b>Iron (FE)</b>                               | Ever     | 151/399          | 1.07 [0.81-1.42] | High     | 74/207           | 0.96 [0.68-1.37] |
|  | Never    | 194/579          | Ref.             | Low      | 77/192           | 1.2 [0.85-1.7]   |
|  |          |                  |                  | No       | 194/579          | Ref.             |
| <b>Lead (PB)</b>                               | Ever     | 169/459          | 1.03 [0.78-1.36] | High     | 80/235           | 0.89 [0.63-1.25] |
|  | Never    | 176/519          | Ref.             | Low      | 89/224           | 1.2 [0.86-1.69]  |
|  |          |                  |                  | No       | 176/519          | Ref.             |
| <b>Low frequency magnetic fields (MF)</b>      | Ever     | 192/536          | 0.96 [0.73-1.27] | High     | 96/268           | 0.96 [0.69-1.34] |
|  | Never    | 153/442          | Ref.             | Low      | 96/268           | 0.97 [0.69-1.36] |
|  |          |                  |                  | No       | 153/442          | Ref.             |
| <b>Methylene chloride (MCH)</b>                | Ever     | 67/196           | 0.92 [0.65-1.3]  | High     | 35/101           | 0.82 [0.51-1.31] |
|  | Never    | 278/782          | Ref.             | Low      | 32/95            | 1.04 [0.65-1.66] |
|  |          |                  |                  | No       | 278/782          | Ref.             |
| <b>Nickel (NI)</b>                             | Ever     | 153/412          | 1.03 [0.78-1.36] | High     | 87/209           | 1.13 [0.81-1.59] |
|  | Never    | 192/566          | Ref.             | Low      | 66/203           | 0.92 [0.65-1.32] |
|  |          |                  |                  | No       | 192/566          | Ref.             |
| <b>Oil mist (OIL)</b>                          | Ever     | 105/324          | 0.87 [0.65-1.17] | High     | 44/162           | 0.68 [0.45-1.03] |
|  | Never    | 240/654          | Ref.             | Low      | 61/162           | 1.06 [0.74-1.53] |
|  |          |                  |                  | No       | 240/654          | Ref.             |
| <b>Other organic solvents (OSOL)</b>           | Ever     | 83/210           | 1.03 [0.74-1.42] | High     | 46/105           | 1.08 [0.71-1.66] |
|  | Never    | 262/768          | Ref.             | Low      | 37/105           | 0.97 [0.62-1.51] |
|  |          |                  |                  | No       | 262/768          | Ref.             |
| <b>Perchloroethylene (PER)</b>                 | Ever     | 5/28             | 0.58 [0.21-1.62] | High     | 2/15             | 0.47 [0.1-2.2]   |
|  | Never    | 340/950          | Ref.             | Low      | 3/13             | 0.7 [0.18-2.76]  |
|  |          |                  |                  | No       | 340/950          | Ref.             |
| <b>Polycyclic aromatic hydrocarbons (PAHs)</b> | Ever     | 123/329          | 1.05 [0.79-1.39] | High     | 59/165           | 0.9 [0.62-1.31]  |
|  | Never    | 222/649          | Ref.             | Low      | 64/164           | 1.21 [0.84-1.74] |
|  |          |                  |                  | No       | 222/649          | Ref.             |
| <b>Toluene (TOLU)</b>                          | Ever     | 72/210           | 0.88 [0.63-1.24] | High     | 47/107           | 1.14 [0.75-1.73] |
|  | Never    | 273/768          | Ref.             | Low      | 25/103           | 0.61 [0.36-1.03] |
|  |          |                  |                  | No       | 273/768          | Ref.             |
| <b>Trichloroethane (TCE)</b>                   | Ever     | 136/358          | 1.16 [0.87-1.54] | High     | 64/185           | 0.98 [0.68-1.42] |
|  | Never    | 209/620          | Ref.             | Low      | 72/173           | 1.35 [0.95-1.93] |
|  |          |                  |                  | No       | 209/620          | Ref.             |
| <b>Trichloroethylene (TRI)</b>                 | Ever     | 115/302          | 1.14 [0.85-1.53] | High     | 59/155           | 1.09 [0.75-1.6]  |
|  | Never    | 230/676          | Ref.             | Low      | 56/147           | 1.2 [0.82-1.75]  |
|  |          |                  |                  | No       | 230/676          | Ref.             |
| <b>Welding fume (WELD)</b>                     | Ever     | 151/399          | 1.07 [0.81-1.42] | High     | 74/202           | 1 [0.7-1.41]     |
|  | Never    | 194/579          | Ref.             | Low      | 77/197           | 1.16 [0.82-1.65] |
|  |          |                  |                  | No       | 194/579          | Ref.             |